## PCT

## WORLD INTELLECTUAL PROPERTY ORGANIZATION International Bureau



### INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification <sup>6</sup>:
C12N 15/29, C07K 14/415, C12N 5/04,
5/14, A01H 5/00, C12N 15/10, 15/82

(11) International Publication Number:

WO 99/19492

(43) International Publication Date:

22 April 1999 (22.04.99)

(21) International Application Number:

PCT/EP98/06977

A2

(22) International Filing Date:

9 October 1998 (09.10.98)

(30) Priority Data:

PO 9745

10 October 1997 (10.10.97) AU

(71) Applicant (for all designated States except US):
RHONE-POULENC AGRO [FR/FR]; 14/20, rue Pierre

Baizet, F-69009 Lyon (FR).

(72) Inventors; and

(75) Inventors/Applicants (for US only): DOUTRIAUX,
Marie-Pascale [FR/FR]; 64, route de Villebon, F-91160
Saulx les Chartreux (FR). BETZNER, Andreas, Stefan
[AU/AU]; 40 Dallachy Place, Page, ACT 2614 (AU).
FREYSSINET, Georges [FR/FR]; 21, rue de Nervieux,
F-69450 Saint Cyr au Mont d'Or (FR). PEREZ, Pascal
[FR/FR]; 17, chemin de la Pradelle, Varennes, F-63450
Chanonat (FR).

(74) Agent: GENIN, Patrick; Rhône-Poulenc Agro, DPI, 14/20, rue Pierre Baizet, F-69009 Lyon (FR). (81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

#### **Published**

Without international search report and to be republished upon receipt of that report.

(54) Title: METHODS FOR OBTAINING PLANT VARIETIES

(57) Abstract

An isolated and purified DNA molecule comprising a polynucleotide sequence encoding a polypeptide functionally involved in the DNA mismatch repair system of a plant.

#### FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
ΑZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav	TM	Turkmenistan
BF	Burkina Faso	GR	Greece		Republic of Macedonia	TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	ΙE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	zw	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's	NZ	New Zealand		
CM	Cameroon		Republic of Korea	PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden	•	
EE	Estonia	LR	Liberia	SG	Singapore		
İ							

10

#### 1

### Methods for Obtaining Plant Varieties

#### TECHNICAL FIELD

The present invention relates to nucleotide sequences which encode polypeptides involved in the DNA mismatch repair systems of plants, and to the polypeptides encoded 5 by those nucleotide sequences. The invention also relates to nucleotide sequences and polypeptide sequences for use in altering the DNA mismatch repair system in plants. The invention also relates to a process for altering the DNA mismatch repair system of a plant cell, to a process for increasing genetic variations in plants and to processes for obtaining plants having a desired characteristic.

#### BACKGROUND OF THE INVENTION

Plant breeding essentially relies on and makes use of genetic variation which occurs naturally within and between members of a family, a genus, a species or a subspecies. Another source of genetic variation is the introduction of genes from other organisms which may or may not be related to the host plant.

Allelic loci or non-allelic genes which constitute or contribute to desired quantitative (e.g. growth performance, yield, etc.) or qualitative (e.g. deposition, content and composition of seed storage products; pathogen resistance genes: etc.) traits that are absent, incomplete or inefficient in a species or subspecies of interest are typically introduced by the plant breeder from other species or subspecies, or *de novo*. This introduction is often done by crossing, provided that the species to be crossed are sexually compatible. Other means of introducing genomes, individual chromosomes or genes into plant cells or plants are well known in the art. They include cell fusion, chemically aided transfection (Schocher et al., 1986, Biotechnology 4: 1093) and ballistic (McCabe et al., 1988, Biotechnology 6: 923), microinjection (Neuhaus et al., 1987, TAG 75: 30), electroporation of protoplasts (Chupeau et al., 1989, Biotechnology 7: 53) or microbial transformation methods such as Agrobacterium mediated transformation (Horsch et al., 1985, Science 227: 1229; Hiei et al., 1996, Biotechnology 14: 745).

However, when a foreign genome, chromosome or gene is introduced into a plant, it will often segregate in subsequent generations from the genome of the recipient plant or plant cell during mitotic and meiotic cell divisions and, in consequence, become lost from the host plant or plant cell into which it had been introduced. Occasionally, however, the introduced genome, chromosome or gene physically combines entirely or in part with the genome, chromosome or gene of the host plant or plant cell in a process which is called recombination.

Recombination involves the exchange of covalent linkages between DNA molecules in regions of identical or similar sequence. It is referred to here as homologous recombination if donor and recipient DNA are identical or nearly identical (at least 99%).

hase sequence identity), and as homeologous recombination if donor and recipient DNA are not identical but are similar (less than 99% base sequence identity).

The ability of two genomes, chromosomes or genes to recombine is known to depend largely on the evolutionary relation between them and thus on the degree of sequence similarity between the two DNA molecules. Whereas homologous recombination is frequently observed during mitosis and meiosis, homeologous recombination is rarely or never seen.

From a breeder's perspective, the limits within which homologous recombination occurs, therefore, define a genetic barrier between species, varieties or lines, in contrast to homeologous recombination which can break this barrier. Homeologous recombination is thus of great importance for plant breeding. Accordingly there is a need for a process for enhancing the frequency of homeologous recombination in plants. In particular, there is a need for a process of increasing homeologous recombination to significantly shorten the length of breeding programs by reducing the number of crosses required to obtain an otherwise rare recombination event.

At least in Escherichia coli, homologous and homeologous recombination are known to share a common pathway that requires among others the proteins RecA, RecB, RecC. RecD and makes use of the SOS induced RuvA and RuvB, respectively. It has been suggested that mating induced recombination follows the Double-Strand Break Repair 20 model (Szostak et al., 1983, Cell 33, 25-35), which is widely used to describe genetic recombination in eukaryotes. Following the alignment of homologous or homeologous DNA double helices the RecA protein mediates an exchange of a single DNA strand from the donor helix to the aligned recipient DNA helix. The incoming strand screens the recipient helix for sequence complementarity, seeking to form a heteroduplex by hydrogen 25 bonding the complementary strand. The displaced homologous or homeologous strand of the recipient helix is guided into the donor helix where it base pairs with its counterpart strand to form a second heteroduplex. The resulting branch point then migrates along the aligned chromosomes thereby elongating and thus stabilising the initial heteroduplexes. Single stranded gaps (if present) are closed by DNA synthesis. The strand cross overs 30 (Holliday junction) are eventually resolved enzymatically to yield the recombination products.

Although in wild type E. coli homologous and homeologous recombination are thus mechanistically similar if not identical, homologous recombination in conjugational crosses E. coli x E. coli occurs five orders of magnitude more frequently than homeologous recombination in conjugational crosses E. coli x S. typhimurium (Matic et al. 1995; Cell 80, 507-515). The imbalance in favour of homologous recombination was shown to be caused largely by the bacterial MisMatch Repair (MMR) system since its

inactivation increased the frequency of homeologous recombination in E. coli up to 1000 fold (Rayssiguier et al. 1989, Nature 342, 396-401).

In E. coli. the MMR system (reviewed by Modrich 1991, Annual Rev Genetics 25, 229-253) is composed of only three proteins known as MutS, MutL and MutH. MutS recognizes and binds to base pair mismatches. MutL then forms a stable complex with mismatch bound MutS. This protein complex now activates the MutH intrinsic single stranded endonuclease which nicks the strand containing the misplaced base and thereby prepares the template for DNA repair enzymes.

During recombination, MMR components inhibit homeologous recombination. In vitro experiments demonstrated that MutS in complex with MutL binds to mismatches at the recombination branch point and physically blocks RecA mediated strand exchange and heteroduplex formation (Worth et al., 1994; PNAS 91, 3238-3241). Interestingly, the SOS dependent RuvAB mediated branch migration is insensitive to MutS/MutL, explaining the observed slight increase in SOS dependent homeologous recombination.

15 Homeologous mating even induces the SOS response, thereby taking advantage of RuvAB induction (Matic et al. 1995, Cell 80, 507-515).

The MMR system thus appears to be a genetic guardian over genome stability in E. coli. In this role it essentially determines the extent to which genetic isolation, that is, speciation, occurs. The diminished sensitivity of the SOS system to MMR, however, allows (within limits) for rapid genomic changes at times of stress, providing the means for fast adaptation to altered environmental conditions and thus contributing to intraspecies genetic variation and species evolution.

The important role of MMR in preserving genomic integrity has been established also in certain eukaryotes. In its efficiency, the human MMR, for example, may even counteract potential gene therapy tools such as triple-helix forming oligonucleotides including RNA-DNA hybrid molecules (Havre et al., 1993, J. Virology 67: 7234-7331; Wang et al., 1995, Mol. Cell. Biol. 15: 1759-1768; Kotani et al., 1996, Mol. Gen. Genetics 250: 626-634; Cole-Strauss et al., 1996, Science 273: 1387-1389). Such oligonucleotides are designed to introduce single base changes into selected DNA target sequences in order to inactivate for example cancer genes or to restore their normal function. The resulting base mismatches however are recognised by the mismatch repair system which then directs removal of the mismatched base, thereby reducing the efficiency of oligonucleotide induced site-specific mutagenesis.

To date, two families of related genes, homologous to the bacterial MutS and MutL genes have been identified or isolated in yeast and mammals (recent reviews by Arnheim and Shibata, 1997, Curr. Opinion Genet. Dev. 7, 364-370; Modrich and Lahue, 1996, Annual Rev. Biochem. 65, 101-133; Umar and Kunkel, 1996, Eur. J. Biochem. 238, 297-307). Biochemical and genetic analysis indicated that eukaryotic MutS homologs (MSH)

25

and MutL homologs (MLH. PMS), respectively, fulfil similar protein functions as their bacterial counterparts. Their relative abundance, however, could reflect different mismatch specificity and/or specialisation for different tissues or organelles or developmental processes such as mitotic versus meiotic recombination.

To date, six different genes homologous to *MutS* have been isolated in yeast (yMSH), and their homologs have been found in mouse (mMSH) and human (hMSH), respectively. Encoded proteins yMSH2, yMSH3 and yMSH6 appear to be the main MutS homologs involved in MMR during mitosis and meiosis in yeast, where the complementary proteins MSH3 and MSH6 alternatively associate with MSH2 to recognise different mismatch substrates (Masischky et al., 1996, Genes Dev. 10, 407-420). Similar protein interactions have been demonstrated for the human homologs hMSH2, hMSH3 and hMSH6 (Acharya et al., 1996, PNAS 93, 13629-13634).

MutL homologs (MLH and PMS), recently reviewed by Modrich and Lahue (1996, Annual Rev. Biochem. 65, 101-133) have so far been found in yeast (yMLH1 and yPMS1), mouse (mPMS2) and human (hMLH1, hPMS1 and hPMS2). The hPMS2 is a member of a family of at least 7 genes (Horii et al., 1994, Biochem. Biophys. Res. Commun. 204, 1257-1264) and its gene product is most closely related to yPMS1. Prolla et al. (1994, Science 265, 1091-1093) presented evidence for yPMS1 and yMLH1 to physically associate with each other and, together, to interact with the MutS homolog yMSH2 to form a ternary complex involved in mismatch substrate binding.

However, while medical interest in mismatch repair has prompted extensive research on MMR in bacteria, yeast and mammals, MMR genes have not been isolated from higher plants prior to the present invention and no attempts to adjust the plant MMR to plant breeding needs have been reported.

#### SUMMARY OF THE INVENTION

According to a first embodiment of the invention, there is provided an isolated and purified DNA molecule comprising a polynucleotide sequence encoding a polypeptide functionally involved in the DNA mismatch repair system of a plant. In one form of this embodiment, the invention provides an isolated and purified DNA molecule comprising a polynucleotide sequence encoding a polypeptide which is homologous to a mismatch repair polypeptide of a yeast or of a human. More particularly, the invention provides polynucleotide sequences encoding polypeptides which are homologous to the mismatch repair polypeptides MSH3 and MSH6 of Saccharomyces cerevisiae. Still more particularly, the invention provides the coding sequences of the genes AtMSH3 and AtMSH6 of Arabidopsis thaliana. as defined hereinbelow, and polynucleotide sequences encoding polypeptides which are homologous to polypeptides encoded by AtMSH3 and AtMSH6.

According to a second embodiment of the invention, there is provided an isolated and purified polypeptide functionally involved in the DNA mismatch repair system of a plant, for example a polypeptide which is homologous to a mismatch repair polypeptide of a yeast or of a human such as a polypeptide encoded by the genes AtMSH3 or AtMSH6 of Arabidopsis thaliana, as defined hereinbelow.

According to a third embodiment of the invention, there is provided an isolated and purified DNA molecule comprising a polynucleotide sequence selected from the group consisting of (i) a sequence encoding a polynucleotide which is capable of interfering with the expression of a plant polynucleotide sequence encoding a polypeptide which is homologous to a mismatch repair polypeptide of a yeast or of a human and thereby disabling said plant polynucleotide sequence; and (ii) a sequence encoding a polypeptide capable of disrupting the DNA mismatch repair system of a plant.

According to a fourth embodiment of the invention there is provided a chimeric gene comprising a DNA sequence selected from the group consisting of (i) a sequence encoding a polynucleotide which is capable of interfering with the expression of a plant polynucleotide sequence encoding a polypeptide which is homologous to a mismatch repair polypeptide of a yeast or of a human and thereby disabling said plant polynucleotide sequence, and (ii) a sequence encoding a polypeptide capable of disrupting the DNA mismatch repair system of a plant: together with at least one regulation element capable of functioning in a plant cell. Examples of such regulation elements include constitutive, inducible, tissue type specific and cell type specific promoters such as 35S, NOS, PR1a, AoPR1 and DMC1. Typically, a chimeric gene of the fourth embodiment will also include at least one terminator sequence, more typically exactly one terminator sequence.

In the third and fourth embodiments, said interference, by said polynucleotide sequence, with the expression of a plant polynucleotide sequence encoding a polypeptide which is homologous to a mismatch repair peptide of a yeast or a human typically occurs by hybridisation or by co-suppression.

According to a fifth embodiment of the invention there is provided a plasmid or vector comprising a chimeric gene of the fourth embodiment. A vector of the fifth 30 embodiment may be, for example, a viral vector or a bacterial vector.

According to a sixth embodiment of the invention, there is provided a plant cell stably transformed, transfected or electroporated with a plasmid or vector of the fifth embodiment.

According to seventh embodiment of the invention, there is provided a plant so comprising a cell of the sixth embodiment.

According to an eighth embodiment of the invention, there is provided a process for at least partially inactivating a DNA mismatch repair system of a plant cell, comprising

transforming or transfecting said plant cell with a DNA sequence of the third embodiment or a chimeric gene of the fourth embodiment or a plasmid or vector of the fifth embodiment, and causing said DNA sequence to express said polynucleotide or said polypeptide.

According to a ninth embodiment of the invention, there is provided a process for increasing genetic variation in a plant comprising obtaining a hybrid plant from a first plant and a second plant, or cells thereof, said first and second plants being genetically different; altering the mismatch repair system in said hybrid plant; permitting said hybrid plant to self-fertilise and produce offspring plants; and screening said offspring plants for plants in which homeologous recombination has occurred. For example, homeologous recombination may be evidenced by new genetic linkage of a desired characteristic trait or of a gene which contributes to a desired characteristic trait.

According to a tenth embodiment of the invention there is provided a process for obtaining a plant having a desired characteristic, comprising altering the mismatch repair system in a plant, cell or plurality of cells of a plant which does not have said desired characteristic, permitting mutations to persist in said cells to produce mutated plant cells, deriving plants from said mutated plant cells, and screening said plants for a plant having said desired characteristic.

In a preferred form of the ninth and tenth embodiments of the invention, the step of altering the mismatch repair system comprises introducing into said hybrid plant, plant, cell or cells a chimeric gene of the fourth embodiment and permitting the chimeric gene to express a polynucleotide which is capable of interfering with the expression of a plant polynucleotide sequence in a mismatch repair gene of the hybrid plant, plant, cell or cells, or a polypeptide capable of disrupting the DNA mismatch repair system of the hybrid plant or cells.

In other embodiments, the invention provides (a) an oligonucleotide capable of hybridising at 45°C under standard PCR conditions to a DNA molecule of the first embodiment; (b) an oligonucleotide capable of hybridising at 45°C under standard PCR conditions to the DNA of SEQ ID NO: 18 and (c) an oligonucleotide capable of hybridising at 45°C under standard PCR conditions to the DNA of SEQ ID NO:30; with the proviso that the oligonucleotide of (a), (b) and (c) is other than SEQ ID NO:1 or SEQ ID NO:2.

#### BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 provides a diagrammatic representation of the primer sequences used to 35 isolate AtMSH3.

Figure 2 is a plasmid map of clone 52, showing restriction enzyme cleavage sites in the 5' half of the full-length cDNA for AtMSH3.

Figure 3 is a plasmid map of clone 13, showing restriction enzyme cleavage sites in the 3' half of the full-length cDNA for AtMSH3.

Figure 4 is a sequence listing of the coding sequence of AtMSH3, together with a deduced sequence of the encoded polypeptide.

Figure 5 is a protein alignment of yeast (Saccharomyces cerevisiae) and Arabidopsis thaliana MSH3 protein.

Figure 6 provides a diagrammatic representation of the primer sequences used to isolate AtMSH6.

Figure 7 is a plasmid map of clone 43, showing restriction enzyme cleavage sites in the 5' half of the full-length cDNA for AtMSH6.

Figure 8 is a plasmid map of clone 62, showing restriction enzyme cleavage sites in the 3' half of the full-length cDNA for AtMSH6.

Figure 9 is a sequence listing of the coding sequence of AtMSH6, together with a deduced sequence of the encoded polypeptide.

Figure 10 is a protein alignment of yeast (Saccharomyces cerevisiae) and Arabidopsis thaliana MSH6 protein.

Figure 11 is a genomic sequence listing of AtMSH6.

Figure 12 is a plasmid map of plasmid pPF13.

Figure 13 is a plasmid map of plasmid pPF14.

Figure 14 is a plasmid map of plasmid pCW186.

Figure 15 is a plasmid map of plasmid pCW187.

Figure 16 is a plasmid map of plasmid pPF66.

Figure 17 is a plasmid map of plasmid pPF57.

Figure 18 is a diagrammatic representation of an antisense gene construction for use in homeologous meiotic recombination.

Figure 19 is a plasmid map of plasmid p3243.

## DETAILED DESCRIPTION OF THE INVENTION

The present invention is based on the inventors' discovery that there exist in higher plants genes which are homologous to MMR genes in E. coli, and to MMR genes in yeasts and humans.

Thus, the inventors have identified genes, herein designated AtMSH3 and AtMSH6, of the plant Arabidopsis thaliana which encode the proteins AtMSH3 and AtMSH6. These plant proteins are homologous to yMSH3 and yMSH6, respectively. The present inventors have isolated cDNAs encoding the proteins AtMSH3 and AtMSH6 and have isolated the complete gene encoding AtMSH6. Given the teaching herein, other genes (for example AtMSH2, and genes of other plants) may be obtained which are involved in DNA mismatch repair in plants, including other genes which encode polypeptides homologous to MMR proteins of yeasts or humans, such as genes which encode

polypeptides homologous to yeast MSH2. MLH1 or PMS2. or to human MLH1. PMS1 or PMS2. For example, given the teaching herein, genes of members of the Brassicaceae family or of other unrelated families, for example the Poaceae, the Solanaceae, the Asteraceae, the Malvaceae, the Fabaceae, the Linaceae, the Canabinaceae, the Dauaceae and the Cucurbitaceae family, and which encode polypeptides homologous to MMR proteins of yeasts or humans may be obtained.

Examples of plants whose genes encoding polypeptides homologous to MMR proteins of yeasts or humans may be obtained given the teaching herein include maize, wheat, oats, barley, rice, tomato, potato, tobacco, capsicum, sunflower, lettuce, artichoke, safflower, cotton, okra, beans of many kinds including soybean, peas, melon, squash, cucumber, oilseed rape, broccoli, cauliflower, cabbage, flax, hemp, hops and carrot.

Within the meaning of the present invention, a first polypeptide is defined as homologous to a second polypeptide if the amino acid sequence of the first polypeptide exhibits a similarity of at least 50% on the polypeptide level to the amino acid sequence of the second polypeptide.

A procedure which may be followed to obtain genes AtMSH3 and AtMSH6 is described in Example 1. Essentially the same technique may be applied to obtain other mismatch repair genes of Arabidopsis thaliana, and essentially the same technique as 20 exemplified herein may be applied to cDNA obtained by reverse transcription of RNA from other plants. Alternatively, given the sequence information disclosed herein, other degenerate oligonucleotide primers, especially oligonucleotides of the invention which are capable of hybridising at 45°C under standard PCR conditions (such as the conditions described in Example 1 using primers UPMU and DOMU) to AtMSH3 and/or AtMSH6 25 may be designed and obtained for use in isolating sequences of plant mismatch repair genes which are homologous to AtMSH3 or AtMSH6, from other plants. Similarly, oligonucleotides of the invention which are capable of hybridising at 45°C under standard PCR conditions to plant mismatch repair genes of plants other than Arabidopsis thaliana also fall within the scope of the present invention and may be utilised to obtain mismatch Typically, such oligonucleotides are capable of 30 repair genes of still other plants. hybridising at 45°C under standard PCR conditions to a DNA molecule which encodes a polypeptide which is homologous to a mismatch repair polypeptide of a yeast or a human. The temperature at which oligonucleotides of the invention hybridise to AtMSH3 and/or AtMSH6, or to plant mismatch repair genes of plants other than Arabidopsis thaliana, or 35 to DNA molecules which encode polypeptides which are homologous to a mismatch repair polypeptide of a yeast or a human may be higher than 45°C, for example at least 50°C, or at least 55°C, or at least 60°C or as high as 65°C.

35

The successful gene isolation disclosed herein demonstrates for the first time the existence of MMR in higher plants and indicates the presence of other plant MMR genes. For example, genes encoding the plant homologs of MSH1. MSH2. MSH4, MSH5, PMS1, PMS2 and MLH1 may be identified given the teaching herein. Such genes, as well as those specifically described herein, separately or in combination, are useful in manipulating the plant MMR for plant breeding purposes. Thus, for example, the plant MMR may be altered by including in a plant cell a polynucleotide sequence as defined herein above with reference to the third embodiment of the invention, and causing the polynucleotide sequence to express either a polynucleotide which disables a plant MMR gene, or a polypeptide which disrupts the plant's MMR system.

The DNA molecule of the third embodiment of the invention includes a polynucleotide sequence (herein referred to as a MMR altering gene) which may for example encode sense, antisense or ribozyme molecules characterised by sufficient base sequence similarity or complementarity to the gene to be altered to permit the antisense or 15 ribozyme molecule to hybridise with the plant MMR gene in vivo or to permit the sense molecule to participate in co-suppression. Alternatively, the MMR altering gene may encode a protein or proteins which interfere with the activity of a plant MMR protein and thus disrupt the plant's MMR system. For example, such encoded proteins may be antibodies or other proteins capable of interfering with MMR protein function, such as by 20 complexing with a protein functionally involved in plant MMR thereby disrupting the MMR of the plant. An example of such a protein is the MSH3 protein of Arabidopsis thaliana described herein or a protein of another plant which is homologous to the MSH3 protein of A. thaliana. For instance, overexpression of MSH3 in a plant cell causes MSH2 present in the cell to be substantially completely complexed, disrupting the 25 mismatch repair mechanism or mechanisms in the cell which are functionally dependent on the presence of a complex of MSH2 with MSH6. Similarly, mismatch repair mechanisms which depend on the presence of a complex of MSH2 and MSH3 may be disrupted by the overexpression of MSH6.

A chimeric gene of the fourth embodiment, incorporating a MMR altering gene, may be prepared by methods which are known in the art. Similarly, the MMR altering gene may be introduced into a plant cell, regenerating tissue or whole plant by techniques known in the art as being suitable for plant transformation, or by crossing. Known transformation techniques include Agrobacterium tumefaciens or A. rhizogenes mediated gene transfer, ballistic and chemical methods, and electroporation of protoplasts.

The MMR altering gene or genes are typically expressed from suitable promoters. Suitable promoters may direct constitutive expression, such as the 35S or the NOS promoter. Usually, however, the promoter will direct either inducible or tissue specific (e.g. callus; embryonic tissue; etc.), cell type specific (e.g. protoplasts; meiocytes; etc.) or developmental (e.g. embryo) expression of the altering gene or genes, in order for the

MMR system to function in tissue types or cell types, or at developmental stages of the plant, in which it is not desirable for the MMR system to be altered. Using such promoters, therefore, the activity of a MMR altering gene may be limited to a specific stage during plant development or it may be altered by controlling conditions external to the plant, and the deleterious effects of a permanently disabled or altered DNA mismatch repair system in a plant may be avoided. Examples of suitable promoters which are not constitutive are known in the art and include inducible promoters such as PR1a (reviewed by Gatz, 1997, Annual Rev. Plant Phys. Plant Mol. Biol. 48: 89), tissue specific promoters such as AoPR1 (Sabahattin et al., 1993, Biotechnology 11: 218), and cell-type specific promoters such as DMC1.

A chimeric gene in accordance with the invention may further be physically linked to one or more selection markers such as genes which confer phenotypic traits such as herbicide resistance, antibiotic resistance or disease resistance, or which confer some other recognisable trait such as male sterility, male fertility, grain size, colour, growth rate, flowering time, ripening time, etc.

The process of the tenth embodiment of the invention provides, for example, a process for generating intraspecies genetic variation by altering the mismatch repair system in a plant cell, in regenerating plant tissue or in a whole plant. The plant cell, regenerating tissue or whole plant includes and expresses one or more MMR altering 20 genes which are capable of altering mismatch repair in the plant cell, regenerating tissue or whole plant. Alteration of MMR may be achieved, for example, by inactivating the genes encoding plant MSH3 and/or plant MSH6. It is preferred to inactivate the plant MSH3 and MSH6 encoding genes at the same time and in the same plant cell, regenerating tissue or whole plant. Typically in this preferred form of the invention 25 inactivation of either plant MSH3 or MSH6 alone is insufficient to substantially alter the plant's mismatch repair system and only when both MSH3 and MSH6 are inactivated simultaneously is the plant's mismatch repair system sufficiently altered to prevent the MMR system from recognising base pair mismatches, base insertions or deletions as a result of DNA replication errors, DNA damage, or oligonucleotide induced site-specific 30 mutagenesis. However, in some applications of the invention, inactivation of only one gene may also be used to cause genomic instability or increase the efficiency of sitespecific mutagenesis.

If desired, the MMR altering gene or genes may later be rendered non-functional or ineffective, or may be removed from the genome of the plant cell, regenerating tissue or whole plant in order to restore mismatch repair in the plant cell, regenerating tissue or whole plant. The MMR altering gene or genes may be inactivated by means of known gene inactivation tools, such as ribozymes, or may be removed from the genome using gene elimination systems known in the art, such as CRE/LOX. It is preferred to render two genes, whose gene products combine to incapacitate MMR, ineffective by separating

the altering genes through segregation. Therefore, in a preferred embodiment of the invention a first plant cell or plant is generated in which only plant MSH3 is incapacitated, and a second plant cell or plant is generated in which only plant MSH6 is incapacitated. The combination of both genomes, for example by crossing, then produces significant MMR deficiency in those cells or plants which have inherited both altering genes. If the altering genes are expressed from unlinked loci, gene segregation restores MMR activity in the progeny of the cells or plants.

In a process of the ninth embodiment of this invention, homeologous recombination is enhanced between different genomes, chromosomes or genes in plant cells or plants by altering MMR in said plant cells or plants. Such genomes, chromosomes or genes are characterised in that they originate from different plant families, genera, species, subspecies, plant varieties or lines. Hybrid plant cells or hybrid plants may be produced by crossing, by cell fusion or by other techniques known in the art. These plant cells or plants are further characterised by expressing one or more genes that are capable of altering mismatch repair in the plant cell or plants.

In the process of the ninth embodiment, the homeologous recombination is typically for the purpose of introducing a desired characteristic in the hybrid plant. In this typical application of the process of the ninth embodiment, and in the process of the tenth embodiment the desired characteristic may be any characteristic which is of value to the plant breeder. Examples of such characteristics are well known in the art and include altered composition or quality of leaf or seed derived storage products (e.g. oil, starch, protein), altered composition or quality of cell walls (e.g. decrease in lignin content), altered growth rate, prolonged flowering, increased plant yield or grain yield, altered plant morphology, resistence to pathogens, tolerance to or improved performance under environmental stresses of various kinds, etc.

In a preferred form of the tenth embodiment, an MMR altering gene is cointroduced along with the homeologous genome, chromosome or gene of another plant
cell or plant into an MMR proficient plant cell or MMR proficient plant to produce a
hybrid plant cell or hybrid plant in which homeologous recombination can occur.

Suitably, the MMR proficient plant cell or MMR proficient plant may also include an
MMR altering gene. For example a gene capable of inactivating plant MSH3 may be cointroduced along with the homeologous genome, chromosome or gene of another plant
cell or plant into an MMR proficient plant cell or MMR proficient plant in which MSH6
is inactivated. A resultant hybrid plant in which homeologous recombination occurs will
include both the MSH3 and MSH6 altering genes and its MMR system will therefore be
inactivated.

In this form of the invention, if hybrid plants are to be produced by crossing, the MMR altering gene preferably originates from the male parent, thus ensuring that the

MMR altering gene is always introduced and is not present in the recipient cell. That is, the MMR of the recipient cell, prior to introduction of the MMR altering gene, is typically proficient. Alternatively, if an MMR altering gene is present in a recipient cell it may be ineffective or inefficient on its own, or it may be linked to an inducible or tissue specific or cell type specific promoter which only renders the MMR altering gene active under limited conditions.

Thus, in a preferred form of the process of the ninth embodiment, the MMR system of the hybrid plant is initially unaltered. In this form of the process, the step of altering the mismatch repair system may comprise introducing into the hybrid plant, or cells thereof, a MMR altering gene, such as by Agrobacterium tumefaciens or A. rhizogenes mediated gene transfer, ballistic and chemical methods, and electroporation of protoplasts.

The MMR altering gene or genes are typically expressed from suitable promoters, as described above. Preferably, the promoter is transcriptionally active in mitotically and meiotically active tissue and/or cells to ensure MMR alteration after chromosome pairing at mitosis and meiosis, respectively. The preferred timing for MMR alteration is at meiosis, because recombinant genomes, chromosomes or genes are directly transmitted to the progeny. A suitable meiocyte specific promoter is for example the *DMC*1 promoter from *Arabidopsis thaliana* ssp. *Ler*. (Klimyuk and Jones, 1997, Plant J. 11, 1-14). However, mitotic homeologous recombination is also a desirable outcome as somatic recombination events can be transmitted to offspring due to the totipotency of plant cells and the lack of predetermined germ cells in plants.

If desired, the MMR altering gene or genes may later be rendered non-functional or ineffective, or may be removed from the hybrid plant or hybrid plant cells, in order to restore mismatch repair in the hybrid plant or hybrid plant cells. The MMR altering gene or genes may be inactivated by means of known gene inactivation tools as described herein above.

#### **EXAMPLES**

## Example 1. Cloning of the AtMSH3 and AtMSH6 coding sequences

Isolation of partial AtMSH3 and AtMSH6 consensus sequences

30

Degenerate oligonucleotides UPMU (SEQ ID NO:1) and DOMU (SEQ ID NO:2)

UPMU CTGGATCCACIGGICCIAA(C/T)ATG

DOMU CTGGATCC(A/G)TA(A/G)TGIGTI(A/G)C(A/G)AA

were used to isolate AtMSH3 and AtMSH6 sequences by PCR amplification.

Primers UPMU and DOMU correspond to conserved amino acid sequences of the proteins MutS (E. coli and S. typhimurium), HexA (S. pneumoniae). Repl (mouse) and Ducl (human). The conserved regions to which they are targeted are TGPNM for UPMU (amino acid positions 852-856 for AtMSH6 and 816-820 for AtMSH3) FATHY or FVTHY

for DOMU (amino acid positions 964-968 for AtMSH6 and 928-932 for AtMSH3, respectively.) These primers have been used to isolate MSH2 and MSH1 from yeast (Reenan and Kolodner, Genetics 132: 963-973 (1992)) and MSH2 from *Xenopus* and mouse (Varlet et al., Nuc. Acids Res. 22:5723-5728 (1994)).

Template single strand cDNA was produced by reverse transcription of 2 µg total RNA from a cell suspension culture of Arabidopsis thaliana ecotype Columbia (Axelos et al. 1989, Mol. Gen. Genetics 219: 106-112). The PCR reaction was performed under the following conditions in a final volume of 100µl: 0.2mM dNTP, 1µM each primer, 1XPCR buffer, lu Taq DNA polymerase (Appligene) in the presence of template cDNA. PCR 10 parameters were 5 minutes at 94°C, followed by 30 cycles of 40 seconds at 95°C, 90 seconds at 45°C, 1 minute at 72°C. The amplification products were cloned into pGEM-T vector (Promega) and sequenced. Two different clones were isolated, S5 (350bp) was homologous to MSH3, S8 (327bp) was homologous to MSH6. Complete cDNA sequences were then isolated according to the Marathon cDNA amplification kit procedure (Clontech). 15 In summary, this procedure involves producing double stranded cDNA by reverse transcription of 2µg polyA+ RNA from the cell suspension culture of Arabidopsis. Adaptors are ligated on each side of the cDNA. The ligated cDNA is used as a template for 5' and 3' RACE PCR reactions in the presence of primers that are specific for the adaptor on one side (AP1 and AP2), and specific for the targeted gene on the other side. A 5' and a 3' 20 fragment that overlap are thus produced for each gene. The complete gene coding sequence can be reconstituted taking advantage of a unique restriction site, if available, in the overlapping region. Specific details of this procedure as it was used to isolate AtMSH3 and AtMSH6 coding regions, are as follows.

## Isolation of AtMSH3 complete coding sequence

25

From the sequence of clone S5, primer 636 (SEQ ID NO:3) was designed:

636 TGCTAGTGCCTCTTGCAAGCTCAT.

Primer AP1 (SEQ ID NO:4) is complementary to a portion of an adaptor sequence which had been ligated to the 5' and 3' ends of *Arabidopsis* cDNA:

API CCATCCTAATACGACTCACTATAGGGC.

PCR performed on the ligated cDNA with primers 636 and AP1 for the 5' RACE PCR was followed by a second round of amplification with the nested primers AP2 (SEQ ID NO:5) and S525 (SEQ ID NO:6)

AP2 ACTCACTATAGGGCTCGAGCGGC

S525 AGGTTCTGATTATGTGTGACGCTTTACTTA

35 (the latter was also designed to correspond to a part of the sequence of clone S5) and produced a 2720bp DNA fragment. Figure 1 provides a diagrammatic representation of the primer sequences used to isolate AtMSH3. Another primer (S51, SEQ ID NO:7)

S51 GGATCGGGTACTGGGTTTTGAGTGTGAGG

was designed closer to the 5' border and permitted the determination of 99bp upstream to the ATG initiation codon. For the 3' RACE PCR, a first PCR reaction was performed with primers AP1 and 635 (SEQ ID NO:8).

- 635 GCACGTGCTTGATGGTGTTTTCAC
- 5 followed by a second round of amplification, using the nested primers AP2 and S523 (SEQ ID NO:9)
- which produced a DNA fragment of 890bp. Both DNA fragments were subcloned into pGEM-T and sequenced. Since PCR amplification using the Expand Long Template PCR System (Boehringer-Mannheim) produced errors in the sequence, new oligonucleotides were designed to isolate those sequences again by PCR, but with the high fidelity DNA polymerase Pfu. PCR with primers 1S5 (SEQ ID NO:10) and S53 (SEQ ID NO:11)
  - 1S5 ATCCCGGGATGGGCAAGCAAAAGCAGCAGACGA
  - S53 GACAAAGAGCGAAATGAGGCCCCTTGG
- amplified the 1244bp fragment clone 52 (SEQ ID NO:12, cloned into pUC18/Sma1). PCR with primers S52 (SEQ ID NO:13) and 2S5 (SEQ ID NO:14)
  - 2S5 ATCCCGGGTCAAAATGAACAAGTTGGTTTTAGTC
  - S52 GCCACATCTGACTGTTCAAGCCCTCGC
- amplified the 2104bp clone 13 (SEQ ID NO:15, cloned into pUC18/Sma1). The complete coding sequence of the AtMSH3 gene was reconstructed in pUC18 by ligating the 5' half of AtMSH3 (clone 52) to the 3' half of AtMSH3 (clone 13) after digesting with BamH1 which has a unique cleavage site in the overlapping region of both clones. This manipulation yielded plasmid pPF26. The Smal fragment from pPF26 contains the complete AtMSH3 coding sequence. The remaining primers referred to in Figure 1 are as follows:
  - S51 GGATCGGGTACTGGGTTTTGAGTGTGAGG (SEQ ID NO:16)
  - S525 AGGTTCTGATTATGTGTGACGCTTTACTTA (SEQ ID NO:17)

Figures 2 and 3 provide plasmid maps of clones 52 and 13 respectively, showing restriction enzyme cleavage sites. The complete AtMSH3 coding sequence (SEQ ID NO:18) 30 is 3246bp long and is shown in Figure 4 together with the deduced sequence (SEQ ID NO:19) of the encoded polypeptide. AtMSH3 is clearly homologous to the yeast and mouse MSH3 genes. A sequence alignment of polypeptides encoded by AtMSH3 and that encoded by Saccharomyces cerevisiae MSH3 is set out in Figure 5.

## Isolation of the AtMSH6 complete coding sequence and genomic sequences

- The same procedure allowed isolation of the AtMSH6 cDNA. Figure 6 provides a diagrammatic representation of the primer sequences used to isolate AtMSH6. For the 5' RACE PCR, primers 638 (SEQ ID NO:20) and AP1 (SEQ ID NO:4)
  - 638 TCTCTACCAGGTGACGAAAAACCG
    allowed the amplification of a 2889 DNA fragment. Primer S81 (SEQ ID NO:21)

15

## S81 CGTCGCCTTTAGCATCCCCTTCCTTCAC

helped define the 142bp upstream to the ATG initiation codon. On the 3' side, RACE PCR was initially performed with primers S823 (SEQ ID NO:22) and AP1 (SEQ ID NO:4),

S823 GCTTGGCGCATCTAATAGAATCATGACAGG

5 and then with the nested primers 637 (SEQ ID NO:23) and AP2 (SEQ ID NO:5).

637 GACAGCGTCAGTTCTTCAGAATGC

to produce a 774bp DNA fragment. As for *AtMSH3*, those fragments were cloned and sequenced. Re-isolation of the DNA sequence using the high fidelity *Pfu* polymerase and newly designed primers 1S8 (SEQ ID NO:24) and S83 (SEQ ID NO:25) (for the 5' side) led to a 2182 bp DNA fragment identified as clone 43 (SEQ ID NO:26, cloned in pUC18/Sma1), and a 1379bp clone identified as clone 62 (SEQ ID NO:27, also cloned in pUC18/Sma1).

1S8 2S8	ATCCCGGGATGCAGCGCCAGAGATCGATTTTGT ATCCCGGGTTATTTGGGAACACAGTAAGAGGATT (SEQ ID
NO:28)	
S82	GCGTTCGATCATCAGCCTCTGTGTTGC (SEQ ID NO:29)
S83	CGCTATCTATGGCTGCTTCGAATTGAG

Figures 7 and 8 provide plasmid maps of clones 43 and 62 respectively, showing restriction enzyme cleavage sites. Clones 43 and 62 were digested by the *Xmn*1 restriction enzyme for which a unique site is present in their overlapping region and then ligated. The complete *AtMSH6* coding sequence (SEQ ID NO:30) is 3330bp long and is shown in Figure 9 together with the deduced sequence (SEQ ID NO:31) of the encoded polypeptide. *AtMSH6* is clearly homologous to the yeast and mouse *MSH6*genes. A sequence alignment of polypeptides encoded by *AtMSH6* and that encoded by *Saccharomyces cerevisiae MSH6* is set out in Figure 10.

An AtMSH6 genomic sequence was also isolated from a genomic DNA library constituted after partial Sau3AI digestion of DNA from the Arabidopsis cell suspension. 8062bp were sequenced that covered the AtMSH6 gene and show colinearity with the cDNA. 16 introns are found scattered along the gene. The complete genomic sequence 30 (SEQ ID NO:98) is shown in Figure 11.

## Example 2. A measure of somatic variation in MMR deficient plants

#### Constructs

Constructs with antisense AtMSH3 or antisense AtMSH6 or both AtMSH3/AtMSH6 under the control of a single 35S promoter have been inserted into the binary vector pPZP121 (Hajdukiewicz et al., Plant Mol. Biol. 23, 793-799) between the right and left borders of the T-DNA. The pPZP121 plasmid confers chloramphenicol resistance to Escherichia coli or Agrobacterium tumefaciens bacteria. The aacC1 gene is carried by the T-DNA and allows selection of transformed plant cells on gentamycin (Hajdukiewicz et al., Plant Mol. Biol. 25, 989-994). For the purpose of expressing antisense constructs. a 35S

promoter/terminator cassette with a polylinker was introduced into pPZP121. The 3' ends of the considered genes have been chosen since this region seems more efficient for antisense inhibition. For AtMSH3 this corresponds to clone 13 (2104bp), for AtMSH6 this corresponds to clone 62 (1379bp). Clone 13 comprises 2104bp of the 3' region that were cut off the pUC18 vector by Sal1/Sst1 restriction, blunted with T4 DNA polymerase and ligated into the T4 DNA polymerase blunted BamHI site of pPZP121/35S, creating clone pPF13. The same procedure was followed for the 3' region of AtMSH6 clone 62 (1379bp) thus creating plasmid pPF14. For the double constructs, the 3' region (from clone 62) of AtMSH6 was introduced ahead of the AtMSH3 region into pPF13 creating pCW186 and reciprocally, the 3' region of AtMSH3 (from clone 13) was introduced ahead of AtMSH6 into pPF14, creating pCW187.

These constructs were introduced into the Arabidopsis cells (as described below) of wildtype Columbia and of the Columbia tester line.

An alternative strategy to antisense inhibition of AtMSH6 comes from experiments of Marra et al. (1998, Proc. Natl. Acad. Sci USA 95, 8568-8573) who show that overexpression of functional MSH3 results in depletion of MSH6 protein in human cells. This depletion may generate a mismatch repair mutant phenotype.

For the purpose of overexpressing functional AtMSH3 protein in plant cells, the complete MSH3 coding region was excised from pPF26 (example 1) by digestion with 20 SmaI, and was inserted into the SmaI site of pCW164. The resulting construct was named pPF66. It contains a complete AtMSH3 gene under the control of the 35S promoter inside the left (LB) and right (RB) border of the T-DNA. This T-DNA also contains the hpt2 gene for gentamycin selection. Plasmid pPF66 was introduced into Arabidopsis cells as described below. One cell clone was selected which clearly overexpressed the AtMSH3 gene as shown by Northern analysis. Figures 12-16 provide plasmid maps of plasmids pPF13, pPF14, pCW186, pCW187 and pPF66, respectively.

#### Construction of tester construct

For the purpose of Forward Mutagenesis Assays, a tester construct was built containing the coding regions for nptII, codA, uidA. All three genes are driven by the 35S promoter and are terminated by the 35S terminator. This construct was obtained by introducing an EcoR1 fragment encoding the codA cassette (2.5kb) and a HindIII fragment encoding the uidA (GUS) cassette (2.4kb) into the pPZP111 vector (Hajdukiewicz et al.,1994, Plant Mol Biol 23: 793-799) which already contained the nptII expression cassette. This new plasmid was named pPF57. NptII is used to select for transformed plant cells. 35 GUS is used to analyse the degree of gene silencing in the construct (i.e. to identify cell lines in which the transgenes are expressed), and codA is used as a marker for forward mutagenesis (described below).

The plasmid map of pPF57 is provided in Figure 17.

#### Plant cell transformation

The constructs are introduced into Agrobacterium by electroporation. Plant cells are then transformed by co-cultivation. A suspension culture of Arabidopsis thaliana cells that 5 has been established by Axelos et al. (1992, Plant Physiol. Biochem. 30, 1-6) may be used. One day old freshly subcultured cells are diluted five times into AT medium (Gamborg B5 medium. 30g/l sucrose, 200µg/l NAA). 10µl of saturated Agrobacterium containing the transforming T-DNA constructs are added to 10ml diluted cells in a 100ml erlenmeyer. The co-cultivation is agitated slowly (80rpm) for 2 days. The cells are then washed 3 times into AT medium and finally resuspended in the same initial volume (10ml). The culture is agitated for 3 days to allow expression before plating on selection plates (AT/BactoAgar 8g/l+gentamycin 50µg/ml). Transformed individual calli are isolated 3 weeks later.

#### Tester Strain

The tester construct on plasmid pPF57 was introduced into Arabidopsis cells of wildtype Columbia using the transformation protocol described above. Among 10 candidate transformants, one cell clone was shown (by Southern analysis) to have a unique T-DNA insertion. All three genes were shown to be functional in this cell line as indicated by resistance to kanamycin, blue staining in the presence of X-Glu (GUS), and sensitivity to 5-fluoro-cytosine (codA).

MMR altering genes (described above) were then introduced individually into the tester line and transformed cells are used for analysis of both Microsatellite Instability and Forward Mutagenesis.

#### Microsatellite analysis

Microsatellites have been described in Arabidopsis (Bell and Ecker, 1994, Genomics 19, 137-144). The present Example is based on a study of instability of microsatellites that are polymorphic amongst different ecotypes. DNA is extracted from the transformed calli. Specific primers have been defined that are used to amplify the microsatellite sequence. One of the two primers is previously P<sup>32</sup> labelled by T4 kinase. In case of a polymorphic variation, new PCR products appear that do not follow the expected pattern of migration on a polyacrylamide gel. This is a commonly observed feature for MMR deficient cells in yeast or mammalian cells.

In particular, the present Example describes a study on microsatellites ca72 (CA<sub>18</sub>), nga172 (GA<sub>29</sub>), and ATHGENEA(A<sub>39</sub>), chosen because they belong to the types predominantly affected in human mismatch repair deficient tumors. The size of these microsatellites is not conserved from one *Arabidopsis* ecotype to the other.

Arabidopsis cells which are transformed with an MMR altering gene (above) and control cells not expressing the MMR altering gene are allowed to form calli. DNA is

rapidly extracted from the calli and is analysed for microsatellite instability as described in detail by Bell and Ecker 1994. Genomics 19, 137-144. In summary, the relevant microsatellite is amplified by PCR using P32 labelled primers. The PCR products are separated on a DNA sequencing gel for size determination. Size differences between microsatellites from transformed and control cells not expressing the MMR altering gene in question indicate microsatellite instability as a result of MMR alteration.

The sequences of primers used for PCR amplification of microsatellites ca72 and nga172 are included in Table 1. PCR amplification of microsatellite ATHGENEA made use of a forward primer containing the sequence

ACCATGCATAGCTTAAACTTCTTG (SEQ ID NO:32)

and of a reverse primer containing the sequence

ACATAACCACAAATAGGGGTGC (SEQ ID NO:33).

The amplification for microsatellite ca72 revealed in Columbia control cells (with respect to the MMR altering gene) a 248 bp long PCR fragment instead of the published length of 124 bp. DNA sequencing verified this fragment as a CA<sub>18</sub> microsatellite.

#### Forward mutagenesis assay

10

Tester cells transformed with antisense AtMSH3 or antisense AtMSH6 or both AtMSH3/AtMSH6 are analysed for the stability of the codA gene. The functional codA gene confers to sensitivity to 5-fluoro-cytosine (5FC), whereas a gene inactivating mutation in codA will confer resistance to 5FC. The frequency of resistant cells is therefore a good indicator of somatic variation as a direct result of MMR alteration. Variants resistant to 5FC are first analysed for GUS activity. If GUS is inactive, 5FC resistance is assumed to be due to gene silencing (all three genes are under the 35S promoter). If GUS is active, 5FC resistance is assumed to be due to forward mutations that have inactivated codA. PCR is then performed on the putative codA mutant genes which is then sequenced to confirm the presence of forward mutations in codA.

Besides codA, other marker genes may also be used for the Forward Mutagenesis Assay such as the ALS gene (conferring sensitivity to valine or to sulfonylurea; Hervieu and Vaucheret, 1996, Mol. Gen. Genet. 251 220-224; Mazur et al. 1987, Plant Physiol. 85 1110-30 1117).

## Example 3. Homeologous meiotic recombination in Arabidopsis thaliana

# A. Construction of a meiocyte specific gene expression cassette comprising the DMC1 promoter and the NOS terminator

(i) The DMC1 promoter may be used as published by Klimyuk and Jones, 1997, 35 Plant J. 11.1-14). To obtain a more convenient alternative for gene cloning, a 3.3 Kb

long subfragment of the *DMC*1 promoter was obtained by PCR from genomic DNA of *Arabidopsis thaliana* (ssp. Landsberg erecta "Ler").

The PCR was done in three rounds:

Round One: A 3.7 Kb long product was obtained using the forward primer 5 DMCIN-A comprising the sequence

## GAAGCGATATTGTTCGTG (SEQ ID NO:34)

and the reverse primer DMCIN-B comprising the sequence

AGATTGCGAGAACATTCC (SEQ ID NO:35).

The weak amplification product was then used as template for round two and three.

o Round Two: A 3.1 Kb long product comprising the promoter and the 5' untranslated leader was obtained using forward primer DMCIN-1, which contained the sequence

## acgcgtcgacTCAGCTATGAGATTACTCGTG (SEQ ID NO:36)

and introduced a Sall cloning site at the 5' end of the promoter fragment, and reverse primer DMCIN-2 which contained the sequence

gctctagaTTTCTCGCTCTAAGACTCTCT (SEQ ID NO:37)

and introduced a XbaI site at the 3' end of the PCR fragment.

Round Three: A 0.2 Kb long product comprising the first exon/intron of the *DMC*1 promoter was obtained using forward primer DMCIN-3, which contained the sequence

gctctagaGCTTCTCTTAAGTAAGTGATTGAT (SEQ ID NO:38)

and introduced a XbaI site at the 5' end of the PCR fragment, and reverse primer DMCIN-4, containing the sequence

tccccgggctcgagagatctccatggTTTCTTCAGCTCTATGAATCC (SEQ ID NO:39) and introduced at the 3' end of the PCR product restriction sites for Ncol, BglII, Xhol and 55 Smal.

The products obtained in round Two and Three were digested with XbaI and subsequently ligated to reconstitute a 3.3 Kb long DMC1 promoter from which the first two in-frame ATG start codons were replaced with a unique restriction site for XbaI. This promoter can be cloned between the restriction sites for SalI and SmaI of p3264, which contains the SacI-EcoRI NOS terminator in pBIN19, to yield the entire expression cassette in pBIN19. This cassette contains the following cloning sites: NcoI, BgIII, XhoI. SmaI and (already present on p3264) KpnI and SacI.

(ii) Another strategy yielded the following convenient DMC1 promoter. A 1.8 kb
 DNA fragment comprising the 3' terminal part of the meiocyte specific DMC1 promoter
 35 was isolated by PCR from purified genomic DNA of Arabidopsis thaliana (ssp. Landsberg erecta "Ler"). The forward PCR primer (DMC1a) contained the sequence

acgcgtcgacGAATTCGCAAGTGGGG (SEQ ID NO:40)

and introduced a Sall cloning site at the 5' end of the promoter fragment. The reverse PCR primer (DMC1b) contained the sequence

tccatggagatctcccgggtacCGATTTGCTTCGAGGG (SEQ ID NO:41) introducing a polylinker region at the 3' end of the promoter fragment. The PCR reaction was carried out with VENT DNA Polymerase (NEB) over 25 cycles using the following cycling protocol: 1 minute at 94°C, 1 minute at 54°C, 2 minutes at 72°C.

The PCR reaction yielded a blunt ended DNA fragment which was digested with restriction enzyme SaII and was cloned into the cleavage sites of restriction enzymes SaII and SmaI in plasmid p2030, a pUC118 derivative containing the SacI-EcoRI NOS terminator fragment of pBIN121. The cloning yielded plasmid p2031, containing the DMC1 promoter-polylinker-NOS terminator expression cassette depicted in Figure 18.

## 10 B. Construction of an MSH3 antisense gene under the control of the DMC1 promoter

A 2.1 kb DNA fragment encoding the carboxyterminal part of AtMSH3 was removed from the polylinker of clone 13 described in Example 1 after (i) digestion with KpnI, (ii) blunting of the DNA ends generated by KpnI and (iii) digestion with BamHI. The isolated fragment was then cloned in antisense orientation downstream of the DMC1 promoter in plasmid p2031, which had been digested with restriction enzymes SmaI and BgIII. This cloning yielded plasmid p2033 (Figure 18).

After digestion of p2033 with EcoRI, a 4.1 kb DNA fragment was recovered comprising the DMC1 promoter, the partial MSH3 cDNA in antisense orientation with respect to the promoter and the NOS terminator. This fragment was cloned into the EcoRI restriction site of plant transformation vector pNOS-Hyg-SCV to yield plasmid p3242 (Figure 18).

## C. Construction of a combined MSH6/MSH3 antisense gene under the control of a single DMC1 promoter

A 3.1 kb fragment, encoding in antisense orientation the partial AtMSH6 and AtMSH3 sequences and the 35S terminator, was isolated from pCW186 by digestion with KpnI. The fragment was treated with Klenow enzyme to blunt both ends. It was then cloned into the SmaI site of plasmid p3243 (a pNOS-Hyg-SCV derivative, illustrated in Figure 19), which had been opened to delete the region between the SmaI sites. Clones containing the fragment in the antisense orientation with respect to the DMC1 promoter (described in A(ii) above) were identified by diagnostic digestion with BamHI. The selected construct was named p3261.

Another practical way of cloning the double antisense gene is as follows. A 1 kb DNA fragment encoding the carboxyterminal part of AtMSH6 is isolated from clone 62 described in Example 1 after digestion of clone 62 plasmid DNA with BamHI, which cleaves in the 5' polylinker region flanking the partial cDNA, and with EcoRI, which cleaves within the cDNA. The isolated fragment is treated with Klenow enzyme to blunt both its ends and is cloned into the recipient plasmid p2033 or p3242. For the purpose of

cloning, the recipient plasmid may be cleaved with either Aval or Ncol and can be blunted with Klenow enzyme to produce blunt acceptor ends for fragment cloning. This cloning yields two opposite orientations of cloned fragment DNA with respect to the DMC1 promoter. These can be identified by diagnostic digestion with restriction enzymes Scal or XmnI in conjunction with Sacl. The selected construct contains the DMC1 promoter, the combined partial cDNAs for AtMSH3 and AtMSH6 (both cloned in antisense orientation with respect to the DMC1 promoter) and the NOS terminator. If the recipient plasmid is p2033, the combined antisense gene under control the single DMC1 promoter is recovered from the vector after EcoRI digestion and is cloned into the EcoRI restriction site of pNOS-Hyg-SCV.

# D. Construction of a full-length MSH3 sense gene under control of the DMC1 promoter for overexpression of functional MSH3 protein

Overexpression of MSH3 protein was shown in human cells (Marra et al., 1998, Proc. Natl. Acad. Sci. USA 95, 8568-8573) to complex all available MSH2 protein. This leaves MSH6 protein without its partner, leading to the degradation of MSH6 protein and eventually to a mismatch repair phenotype.

This phenomenon is exploited to increase homeologous meiotic recombination in Arabidopsis as an alternative to antisense inhibition of AtMSH6. For this purpose the full-length cDNA encoding AtMSH3 is isolated from plasmid pPF66 by digestion with SmaI and is cloned into the SmaI site of the DMC1 expression cassettes described in A(i).

# E. Selection of Recombination markers on homeologous chromosomes of Arabidopsis thaliana subspecies Landsberg erecta (Ler), Columbia (Col) and C24, respectively

## E(i). Visual recombination markers in Arabidopsis th. subspecies C24:

Agrobacterium mediated transformation with a T-DNA containing a 35S-GUS gene, inactivated by insertion of a 35S-Ac transposable element (Finnegan et al., 1993, Plant Mol. Biol. 22, 625-633), had yielded a C24 line in which the T-DNA construct was integrated into chromosome 2. Genetic and molecular analysis of this line shows that the Ac transposon had excised from its T-DNA locus thereby restoring GUS activity, but had re-inserted into the chromosome at a distance of 16.4 cM, where it stayed fixed (due to disablement of Ac) within the chlorina gene. Insertional inactivation of the chlorina gene caused a bleached phenotype in those plants that were homozygous for this mutation. Because of the two linked phenotypic markers, chlorina3A:Ac and GUS, this C24 line was used in crosses to wildtype Ler for analysis of meiotic homeologous recombination on chromosome 2 in conjunction with molecular recombination markers.

### 35 E(ii). Visual recombination markers in Arabidopsis th. Ler:

The Ler line NW1 (obtained from NASC, Nottingham, UK) contains one recessive visual marker per chromosome. i.e. an-1 on Chr.1, py-1 on Chr.2, gl1-1 on Chr.3, cer2-1

on Chr.4, and msl-1 on Chr.5. This line is used in crosses to wildtype C24 which expresses an MMR altering gene for analysis of meiotic homeologous recombination on chromosomes 1-5 in conjunction with molecular recombination markers listed in Table 1.

Other Ler lines from NASC have several visual markers in close proximity to each other on the same chromosome. When these lines are used for hybrid production, analysis of homeologous meiotic recombination may make use entirely of visual recombination markers. Particularly suitable for crossing to C24 wildtype that is expressing a MMR altering gene are the following Ler lines:

NW22: relative markers are dis1 - (4 cM) - ga4 - (11 cM) - th1 on chromosome 1.

NW10: relevant markers are tz-201 - (5 cM) - cer3 on chromosome 5.

NW134, relevant markers are ttg - (4 cM) - ga3 on chromosome 5.

10

NW24 (abi3-1) and NW64 (gl1-1). When present in the same plant on chromosome 3, abi3-1 and gl1-1 are 13 cM apart. Since this marker combination is not available from NASC, we have combined these markers by crossing line NW24 to line NW64. The F1 offspring were allowed to self-fertilise and to produce F2 seeds. F2 Plants which carry both markers as homozygous traits on the same chromosome can be identified firstly, by germinating F2 seeds on germination medium containing selective concentrations of abscisic acid, and subsequently, by identifying among the abscisic acid resistant plants those individuals which show the glabra phenotype.

## 20 E(iii) Molecular recombination markers in Col, Ler and C24:

The genome of Arabidopsis thaliana is interspersed with unique base sequences arranged as simple tandem repeats. Allelic repeats can vary in length between different Arabidopsis subspecies and when amplified by PCR yield diagnostic DNA products of different length named Simple Sequence Length Polymorphisms (SSLPs). Many SSLPs have been genetically mapped and have been assigned to unique chromosome locations on the recombinant inbred map (Bell and Ecker, 1994, Genomics 19, 137-144; Lister and Deans lines, Weeds World 4i, May 1997).

In Table 1 are listed 28 mapped and established SSLPs between Ler and Col. A number of PCR primer pairs are described herein (SEQ ID NO:42 to SEQ ID NO:97) which also yielded SSLPs between C24 and Ler (19 SSLPs) or between C24 and Col (25 SSLPs), respectively. Polymorphic SSLPs can be used as molecular markers in the analysis of homeologous recombination between genomes from these subspecies.

The PCR reactions (25 µL) were carried out over 35 cycles (15 seconds at 94°C, 30 seconds at 55°C and 30 seconds at 72°C), with 0.25 U Taq DNA polymerase and 0.6 µg genomic DNA in reaction buffer containing 2 mM MgCl<sub>2</sub>. PCR products were separated by agarose gel electrophoresis (4% ultra high resolution agarose) and visualised by ethidiumbromide staining. The results from the PCR experiments are summarised in

Table 1, which also shows the sequence of PCR primers, primer annealing temperature (Tm), PCR product length and chromosome location of SSLP (with respect to the RI map of May 1997, Weeds World 4i).

### F. Production of hybrid plants

C24 plants heterozygous for chlorina3A:Ac/GUS are crossed as male to emasculated wildtype Ler to produce Ler/C24(chlorina3A, GUS) hybrid seeds.

Due to the heterozygosity of the C24 parent, only 50 % of hybrid plants have inherited the chlorina3A:Ac/GUS locus. The remaining 50% of hybrid plants are wildtype with respect to chlorina3A:Ac/GUS. Since the mutant locus is linked to a kanamycin resistance gene (contained on the same T-DNA as GUS) mutant plants can be pre-selected by germinating hybrid seeds on germination medium containing 50 mg/L kanamycin.

Ler plants homozygous for the five chromosome markers are male sterile (ms1-1) and are crossed without emasculation to wildtype C24 to produce Ler(an-1, py-1, gl1-1, cer2-1, ms1-1)/C24 hybrid seeds.

Other Ler plants, which are male fertile, are crossed after emasculation of the female parent to produce Ler/C24 hybrid seeds.

G. Introduction of MSH3 and MSH6/3 antisense genes into Arabidopsis and analysis of meiotic homeologous recombination

## (i) Transformation of hybrid plants and analysis of homeologous meiotic recombination

The plant transformation vectors comprising the antisense genes described in (B) and (C) above are introduced into Agrobacterium tumefaciens strain AGL1 (Lazo et al., 1991, Bio/Technology 9, 963-967) by electroporation. Recombinant Agrobacterium clones are selected on LB medium containing 50 mg/L rifampicin and 100 mg/L carbenicillin. Selected clones are used to infect roots of Arabidopsis hybrid plants (described in (F) above) using the root transformation protocol of Valvekens et al. (1988, PNAS 85, 5536-5540) except that shoot and root inducing media contain hygromycin (10 mg/L) instead of kanamycin.

Plants regenerated from roots of hybrid plants are genetic clones of root donating plants and therefore are again genetic hybrids of two Arabidopsis subspecies described in 30 (F). However, in contrast to the root donating plants, the regenerated hybrid plants also contain the introduced transgene and the co-introduced hygromycin resistance gene with the latter allowing these plants to grow on hygromycin containing culture medium.

Hygromycin resistant plants are then allowed to enter the reproductive phase and to produce gametes by meiotic divisions of microspore and megaspore mothercells. At meiosis, the *DMC*1 promoter is activated and can direct the expression of antisense genes described in (B) and (C) above, leading to decreased *MSH*6 and/or *MSH*3 gene

expression. This in turn depletes the gamete mothercells of MSH6 and/or MSH3 protein, thus causing alteration of MMR during meiotic divisions and increasing the recombination frequency between homeologous chromosomes.

Transgenic plants are then allowed to self-fertilise and to produce seeds. These seeds (F2 seeds with respect to hybrid production), and the plants derived therefrom, carry the homeologous recombination events which can be identified by using the visual and molecular recombination markers described in (E) above.

In case of homeologous recombination between chromosomes of Ler and C24(chlorina3A:Ac, GUS), the analysis concentrates on chromosome 2 by selecting plants showing the visual phenotypic marker chlorina. This marker thus serves as a reference point as it indicates that respective chromosomes 2 originate from C24. Other markers, such as GUS or molecular markers, on chromosome 2 may then be used to identify chromosomal regions which are derived from the Ler chromosome as a result of homeologous recombination. F2 plants of control transformants not expressing the antisense gene(s) can be analysed in parallel and the results can be used for comparison to homeologous recombination results obtained in antisense plants.

# (ii) <u>Transformation of C24 wildtype</u>, hybrid plant production and analysis of homeologous meiotic recombination

Introduction of MMR altering genes into wildtype C24 is done using the root transformation protocol as described in G(i) for transformation of hybrid plants. Transformed plants are selected by resistance to either 10 mg/L hygromycin (in case of transformation with T-DNA's derived from pNOS-Hyg-SCV) or to 50 mg/L kanamycin (in case of transformation with T-DNA's derived from pBIN19).

Transgenic plants are then allowed to self-fertilise and to produce seeds (T1 seeds).

Segregation of the antibiotic resistance gene in the T1 population then indicates the number of transgene loci. Lines with a single transgene locus (indicated by a 3:1 ratio of resistant:sensitive plants) are selected and are bred to homozygosity. This is done by collecting selfed seeds (T2) from T1 plants and subsequent testing of at least four independent T2 seed populations for segregation of the antibiotic resistance gene. T2 populations which do not segregate the antibiotic resistance gene are assumed to be homozygous for both the resistance gene and the linked MMR altering gene.

C24 plants homozygous for the MMR altering gene are then crossed to Ler lines homozygous for recessive visual markers (see E(ii)) to produce C24/Ler hybrid plants as described in (F). These F1 hybrids are then allowed to enter the reproductive phase and to produce gametes by meiotic division of microspore and megaspore mothercells. At meiosis, the DMC 1 promoter is activated and can direct the expression of antisense or sense genes described in (B), (C) and (D) above, leading to decreased MSH6 and/or MSH3 gene expression. This in turn depletes the gamete mothercells of MSH6 and/or MSH3

protein, thus causing alteration of MMR during meiotic divisions and increasing the recombination frequency between the homeologous chromosomes of C24 and Ler. Recombination events are then scored in the F2 generation.

For recombination analysis, the hybrid plants are allowed to self-fertilise and to produce F2 seeds. F2 plants are then preselected for a first visual marker. Since this marker is recessive, its visual presence indicates homozygosity for Ler DNA at the relevant locus. Those F2 plants which show this first visual marker are then analysed for the presence or absence of a second visual marker which in the Ler parent is closely linked to the first marker. Absence of the second visual marker indicates recombination between the relevant C24 and Ler chromosomes between the first and second marker. The frequency of recombination in transgenic hybrids is compared to the recombination frequency in control hybrids not expressing the MMR altering gene.

Examples of recombination analysis are the following.

The Ler line NW22(dis1, ga4, th1) is used for crosses to transformed C24.

F2 plants are preselected first for thiamine requirement (th1) and then are further analysed for re-appearance of wildtype height (loss of ga4) and/or re-appearance of normal trichomes (loss of dis1) as a result of recombination.

The Ler line NW10(1z-201, cer3) is used for crosses to transformed C24.

F2 plants are then preselected first for thiazole requirement (tz) and then are further 20 analysed for re-appearance of normal, i.e. non-shiny stems (loss of cer3) as a result of recombination.

The Ler line NW134 (ttg, ga3) is used for crosses to transformed C24. F2 plants are first preselected for dwarfish appearance (ga3) and are then analysed for re-appearance of trichomes (loss of ttg) as a result of recombination.

Ler plants homozygous for abi3-1 and gl1-1 are used for crosses to transformed C24. F2 plants are first preselected for their ability to germinate in the presence of abscisic acid and are then analysed for re-appearance of trichomes on the leaves (loss of gl1-1) as a result of recombination.

In the case of homeologous recombination between transformed C24 and the Ler line NW1 (an-1, py-1, gl1-1, cer2-1, ms1-1), recombination analysis is similar the one described above, except that the second marker is not a visual marker but has to be a molecular marker. This is because the Ler parent carries only one visual marker per chromosome.

	L	TABLE 1: SS	SSLP Markers in Arabidopsis thaliana Subspecies	aliana S	ubspecies	-	
Chromosome	RI Map Position	PCR Primer Pair	Primer Sequence	Tm [°C]	length/COL	length/LER	length/C24
	2.3	ATEAT! F	GCCACTGCGTGAATGATATG	57.8	172	162	162
		ATEAT! R	CGAACAGCCAACATTAATTCCC	58.2			
_	9.3	NGA63 F	AACCAAGGCACAGAAGCG	57.3	Ξ	68	120
		NGA63 R	ACCCAAGTGATCGCCACC	59.6			
	40.1	NGA248 F	TACCGAACCAAAACACAAAGG	56.1	143	129	no amplific.
		NGA248 R	TCTGTATCTCGGTGAATTCTCC	58.2			
	81.2	NGA128 F	GGTCTGTTGATGTCGTAAGTCG	60.1	180	061	no amplific.
		NGA128 R	ATCTTGAAACCTTTAGGGAGGG	58.2			
-	81.2	NGA280 F	CTGATCTCACGGACAATAGTGC	60.1	105	85	82
		NGA280 R	GGCTCCATAAAAGTGCACC	57.8			
	111.4	NGA111 F	CTCCAGTTGGAAGCTAAAGGG	39	128	162	170
		NGA111 R	TGTTTTTAGGACAAATGGCG	70			
=	ca. 7.5	NGA168 F	CCTTCACATCCAAAACCCAC	57.8	213	217	208
		NGA 168 R	GCACATACCCACAACCAGAA	57.8			

Ca. 48   NGA1126L   CGCTACGCTTTTCGGTAAAG   57.8   191   199   1								
Ca. 19   NGA1126R   GCACGTCCAAGTCAACC   59.9		97 63	NGA11261	CGCTACGCTTTTCGGTAAAG	57.8	161	661	961
62.2   NGA361L   AAAGAGATGAGAATTTGGAC   51.7   114   120   11   120		CA. 40	NGA 1126R	GCACAGTCCAAGTCACAACC	59.9			
62.2   NGA361L   AAAGAGATTGGAC   51.7   114   120   114   120								
NGA168 F TCGTCTACTGCACTGGGG		62.2	NGA361L	AAAGAGATGAGAATTTGGAC	51.7	114	120	114
73   NGA168 F   TCGTCTACTGCACTGCG   59,6   151   135   151   135   152   153   153   153   154   154   155			NGA361R	ACATATCAATATATTAAAGTAGC	49.5			
13   NGA168 F   TCGTCTACTGCACTGCG   59,6   151   135     NGA168 R   GAGGACATGTATAGGAGCCTCG   61.9   141   209     Ca. 77   AIBBIO2 L   TGACCTCCTCTCCATGGAG   59.9   141   209   141     Ca. 83   AIBBIO2 R   TTAACAGAAACCTATC   54.5   148   148     Ca. 83   AIBBIO2 R   TTAACAGAAACCTATTCATTG   54.1   146   148     I 3.4   NGA12 F   AGCCAATGTCCATTTCTCC   57.8   196   147     II 12.8   NGA12 F   GAAAAACGCTACTTCTTGGG   56.1   119   147     II 12.8   NGA12 F   GAAAAACGCTACTTTCTGGG   56.1   119   147     II 17.5   NGA12 F   CATGCAATTTCATGGG   58.2   107   89     II 17.5   NGA162 F   CATGCAATTTCCTTGGG   55.8   107   89     II 17.5   NGA162 R   CTCTGTCACTTTTCCTGGG   60.1   100   100     II 17.5   NGA162 R   CATGCAATTTCCTTTGGG   60.1   100   100     II 17.5   NGA162 R   CTCTGTCACTTTTCCTTGG   60.1   100   100     II 17.5   NGA162 R   CTCTGTCACTTTTCCTTGG   60.1   100   100     II 17.5   NGA162 R   CTCTGTCACTTTTTCCTTGG   60.1   100   100     II 17.5   NGA162 R   CTCTGTCACTTTTCCTTGG   60.1   100   100     II 17.5   NGA162 R   CTCTGTCACTTTTCCTTGG   60.1   100   100     II 17.5   NGA162 R   CTCTGTCACTTTTTCCTTGG   60.1   100   100   100     II 17.5   NGA162 R   CTCTGTCACTTTTTCCTTGG   60.1   100								
Ca. 77   Alibio2 L   TGACCTCCTTTCCATGGAG   59.9   141   209   141   209   141   209   141   209   141   209   141   209   141   209   141   209   141   209   141   209   141   209   142   209   20   20   20   20   20   20		73	A168	TCGTCTACTGCACTGCCG	9.69	151	135	135
ca. 77         AnbBIO2 L         TGACCTCCTCTTCCATGGAG         59.9         141         209           ca. 83         AnbBIO2 R         TTAACAGAAACCCAAAGCTTTC         54.1         146         148           ca. 83         AnhUBIQUE L         AGGCAAATGTCCATTTCATTG         54.1         146         148           1         a. 83         AnhUBIQUE R         ACGACATGGCAGATTTCTCC         57.8         148           1         3.4         NGA172 F         AGCTGCTTCCTTATAGCGTCC         60         162         136           1         12.8         NGA172 F         AGCTGCTTCCTTATAGCGTCC         55.4         147           1         12.8         NGA126 F         GAAAAAACGCTACTTTCGTGG         55.4         147           1         12.8         NGA126 R         CAAGAGCAATTTCGTGGG         56.1         119         147           1         17.5         NGA126 R         CAAGAGCAATTTCCTTGAGG         55.8         107         89           II         17.5         NGA162 R         CATGCAATTTCCTTGAGG         55.8         107         89           II         17.5         NGA162 R         CTCTGTCACTTTTTCCTTGG         60.1         107         89			A168	GAGGACATGTATAGGAGCCTCG	61.9			
ca. 77         AthBIO2 L         TGACCTCCTCTTCCATGGAG         59.9         141         209           ca. 83         AthUBIQUE L         AGGCAAATGTCCATTTCATTG         54.1         146         148           1         ca. 83         AthUBIQUE R         AGGCAATGGCAGATTTCATTG         57.8         148           1         3.4         NGA172 F         AGCTGCTTCCTTATAGCGTCC         60         162         136           1         12.8         NGA172 R         CATCCGAATGCCATTGTTC         55.4         147         147           1         12.8         NGA126 F         GAAAAAACGCTACTTTCGTGG         56.1         119         147           1         12.8         NGA126 F         GAAAAAACGCTACTTTCGTGG         58.2         107         89           1         17.5         NGA162 F         CATGCATTTCCTTGAGG         55.8         107         89           1         17.5         NGA162 F         CATGCATTTCCTTTTCGTGGG         55.8         107         89           1         17.5         NGA162 R         CTCTGTCATTTTCCTTGGG         55.8         107         89								
AthBIO2 R   TTAACAGAAACCCAAAGCTTTC   54.5		ca. 77	AthBIO2 L	TGACCTCCTCTTCCATGGAG	59.9	141	209	139
ca. 83         AhhUBIQUE L         AGGCAAATGTCCATTTCATTG         54.1         146         148           1         3.4         AthUBIQUE R         ACGACATGGCAGATTTCTCC         57.8         136           1         3.4         NGA172 F         AGCTGCTTCCTTATAGCGTCC         60         162         136           1         12.8         NGA172 R         CATCCGAATGCCATTGTTC         55.4         147           1         12.8         NGA126 F         GAAAAAACGCTACTTTCGTGG         56.1         119         147           1         12.8         NGA126 F         GAAAAAACGCTACTTTCGTGG         58.1         109         89           1         17.5         NGA162 F         CATGCATTTGCATCTGAGG         55.8         107         89           1         17.5         NGA162 F         CATGCATTTTCCTTGAGG         60.1         60.1         60.1         60.1			AthBIO2 R	TTAACAGAAACCCAAAGCTTTC	54.5			
ca. 83         AthUBIQUE L         AGGCAAATGTCCATTTCATG         54.1         146         148           1         AthUBIQUE R         ACGACATGCCAGATTTCTCC         57.8         1         1           1         3.4         NGA172 F         AGCTGCTTCCTTATAGCGTCC         60         162         136           1         12.8         NGA172 R         CATCCGAATGCCATTGTTC         55.4         1         147           1         12.8         NGA126 F         GAAAAAAACGCTACTTTCGTGG         56.1         119         147           1         12.8         NGA126 F         CAAGAGCAATATCAAGAGCAGC         58.2         107         89           1         17.5         NGA162 F         CATGCAATTTGCATCTGAGG         55.8         107         89           1         17.5         NGA162 F         CATGCAATTTGCATCTGAGG         55.8         107         89           1         17.5         NGA162 F         CATGCTCTTTTCCTGTGG         60.1         60.1         80.1								
AthUBIQUE R ACGACATGGCAGATTTCTCC   57.8			AthUBIQUE L	AGGCAAATGTCCATTTCATTG	54.1	146	148	148
3.4         NGA172 F         AGCTGCTTCCTTATAGCGTCC         60         162         136           12.8         NGA172 R         CATCCGAATGCCATTGTTC         55.4         119         147           12.8         NGA126 F         GAAAAAACGCTACTTTCGTGG         56.1         119         147           NGA126 R         CAAGAGCAATATCAAGAGCAGC         58.2         107         89           17.5         NGA162 F         CATGCATTTGCATCTGAGG         55.8         107         89           NGA162 R         CTCTGTCACTCTTTTCCTTGG         60.1         80.1         107         89			AthUBIQUE R	ACGACATGCCAGATTTCTCC	57.8			
3.4         NGA172 F         AGCTGCTTCCTTATAGCGTCC         60         162         136           12.8         NGA172 R         CATCCGAATGCCATTGTTC         55.4         119         147           12.8         NGA126 F         GAAAAAACGCTACTTTCGTGG         56.1         119         147           NGA126 R         CAAGAGCAATATCAAGAGCAGC         58.2         109         89           17.5         NGA162 F         CATGCAATTTGCATCTGAGG         55.8         107         89           NGA162 R         CTCTGTCACTCTTTTCCTCTGG         60.1         80.1         107         89								
NGA172 R   CATCCGAATGCCATTGTTC   55.4		3.4	A172	AGCTGCTTCCTTATAGCGTCC	09	162	136	140
12.8         NGA126 F         GAAAAAACGCTACTTTCGTGG         56.1         119         147           NGA126 R         CAAGAGCAATATCAAGAGCAGC         58.2         2         2         3         3         3         3         3         3         3         4 <t< th=""><th></th><th></th><td>A172</td><td>CATCCGAATGCCATTGTTC</td><td>55.4</td><td></td><td></td><td></td></t<>			A172	CATCCGAATGCCATTGTTC	55.4			
12.8         NGA126 F         GAAAAACGCTACTTTCGTGG         56.1         119         147           NGA126 R         CAAGAGCAATATCAAGAGCAGC         58.2   17.5         NGA162 F         CATGCAATTTGCATCTGAGG         55.8         107         89           NGA162 R         CTCTGTCACTCTTTCCTCTGG         60.1								
NGA126 R   CAAGAGCAATATCAAGAGCAGC		12.8		GAAAAACGCTACTTTCGTGG	56.1	119	147	no amplific.
17.5 NGA162 F CATGCAATTTGCATCTGAGG 55.8 107 89 NGA162 R CTCTGTCACTCTTTCCTCTGG 60.1			A126	CAAGAGCAATATCAAGAGCAGC	58.2			
17.5         NGA162 F         CATGCAATTTGCATCTGAGG         55.8         107         89           NGA162 R         CTCTGTCACTCTTTCCTCTGG         60.1 <th></th> <th></th> <th></th> <th></th> <th></th> <th></th> <th></th> <th></th>								
CTCTGTCACTCTTTTCCTCTGG	=	17.5	NGA162 F	CATGCAATTTGCATCTGAGG	55.8	107	68	no amplific.
			NGA162 R	CTCTGTCACTCTTTTCCTCTGG	1.09			

	81.8	NGA6 F	TGGATTTCTTCCTCTTCAC	56.1	143	123	143
		NGA6 R	ATGGAGAGCTTACACTGATC	56.1			
ΛI	19.8	NGA12 F	AATGTTGTCCTCCCTCCTC	59.9	247	234	220
		NGA12 R	TGATGCTCTCTGAAACAAGAGC	58.2			
\\	24.1	NGA8 F	GAGGCAAATCTTTATTTCGG	56.1	154	861	061
		NGA8 R	TGGCTTTCGTTTATAAACATCC	54.5			
<u>&gt;1</u>	102	NGA1107 L	GCGAAAAACAAAAAATCCA	50.2	150	140	140
		NGA1107 R	CGACGAATCGACAGAATTAGG	58			
>	11.8	NGA225 F	GAAATCCAAATCCCAGAGAGG	58	611	189	119
		NGA225 R	TCTCCCCACTAGTTTTGTGTCC	60.1			
>	16.7	NGA249 F	TACCGTCAATTTCATCGCC	55.4	125	115	115
		NGA249 R	GGATCCCTAACTGTAAAATCCC	58.2			
>	19.9	CA72 F	AATCCCAGTAACCAAACACACA	56.3	124	110	110
		CA72 R	CCCAGTCTAACCACGACCAC	61.9			
>	20	NGA151 F	GITTTGGGAAGITTTGCTGG	55.8	150	120	130
		NGA151 R	CAGTCTAAAAGCGAGAGTATGATG	58.6			

						.03	170
>	24	NGA106 F	GTTATGGAGITTICTAGGGCACG	1.09	157	(2)	00.1
		NGA106 R	TGCCCCATTTGTTCTTCTC	55.8			
^	37.8	NGA139 F	AGAGCTACCAGATCCGATGG	59.9	174	132	132
-		NGA139 R	GGTTTCGTTTCACTATCCAGG	55.8			
							-
>	50	NGA76 F	GGAGAAATGTCACTCTCCACC	1.09	231	> 250	300
		NGA76 R	AGGCATGGGAGACATTTACG	57.8			
>	61.1	ATHSO191 L	CTCCACCAATCATGCAAATG	55.8	148	156	146
•		ATHSO191 R	TGATGTTGATGGAGATGGTCA	53.7			
						3	
>	81.7	NGA129 F	TCAGGAGGAACTAAAGTGAGGG	1.09	177	6/.1	7/1
		NGA129 R	CACACTGAAGATGGTCTTGAGG	60.1			

#### **CLAIMS**

- 1. An isolated and purified DNA molecule comprising a polynucleotide sequence encoding a polypeptide functionally involved in the DNA mismatch repair system of a plant.
- 2. A DNA molecule according to claim 1 wherein said polypeptide is homologous to a mismatch repair polypeptide of a yeast or of a human.
- 3. A DNA molecule according to claim 1 wherein said polypeptide is homologous to AtMSH3 (SEQ ID NO: 19) or to AtMSH6 (SEQ ID NO: 31).
- 4. An isolated and purified polypeptide functionally involved in the DNA nismatch repair system of a plant.
  - 5. A polypeptide according to claim 4 which is homologous to a mismatch repair polypeptide of a yeast or of a human.
- 6. An isolated and purified polypeptide selected from the group consisting of a polypeptide encoded by the gene AtMSH3 (SEQ ID NO: 18), a polypeptide encoded by the gene AtMSH6 (SEQ ID NO:30), polypeptides homologous to a polypeptide encoded by the gene AtMSH3 (SEQ ID NO: 18) and polypeptides homologous to a polypeptide encoded by the gene AtMSH6 (SEQ ID NO:30).
- 7. An isolated and purified DNA molecule comprising a polynucleotide sequence selected from the group consisting of (i) a sequence encoding a polynucleotide which is capable of interfering with the expression of a plant polynucleotide sequence encoding a polypeptide which is homologous to a mismatch repair polypeptide of a yeast or of a human and thereby disabling said plant polynucleotide sequence; and (ii) a sequence encoding a polypeptide capable of disrupting the DNA mismatch repair system of a plant.
- 8. A DNA molecule according to claim 7 comprising a polynucleotide sequence 25 encoding a polynucleotide capable of interfering with the expression of a plant polynucleotide sequence encoding a polypeptide which is homologous to a mismatch repair polypeptide of a yeast or of a human and thereby disabling said plant polynucleotide sequence.
- 9. A DNA molecule according to claim 8 wherein said polynucleotide is capable 30 of interfering with the expression of a plant polynucleotide sequence is a sense polynucleotide, an antisense polynucleotide or a ribozyme.
  - 10. A DNA molecule according to claim 7 comprising a polynucleotide sequence encoding a polypeptide capable of disrupting the DNA mismatch repair system of a plant.

- 11. A DNA molecule according to claim 10 wherein said polypeptide is homologous to AtMSH3 (SEQ ID NO: 19) or to AtMSH6 (SEQ ID NO: 31).
- 12. A DNA molecule according to claim 10 further comprising a regulation element capable of causing overexpression of said polypeptide in a cell of said plant.

### 5 13. A chimeric gene comprising:

a DNA sequence selected from the group consisting of (i) a sequence encoding a polynucleotide capable of interfering with the expression of a plant polynucleotide sequence encoding a polypeptide which is homologous to a mismatch repair polypeptide of a yeast or of a human and thereby disabling said plant polynucleotide sequence, and (ii) a sequence encoding a polypeptide capable of disrupting the DNA mismatch repair system of a plant; and

at least one regulation element capable of functioning in a plant cell.

- 14. A chimeric gene according to claim 13 wherein said regulation element is selected from constitutive, inducible, tissue type specific and cell type specific promoters.
- 15. A chimeric gene according to claim 13 comprising a DNA sequence encoding a polypeptide capable of disrupting the DNA mismatch repair system of a plant, wherein said regulation element is capable of causing overexpression of said polypeptide in a cell of said plant.
- 16. A chimeric gene according to claim 13 wherein said regulation element is selected from the group consisting of 35S, NOS, PR1a, AoPR1 and DMC1.
  - 17. A plasmid or vector comprising a chimeric gene according to any one of claims 13-16.
  - 18. A plant cell stably transformed, transfected or electroporated with a plasmid or vector according to claim 17.
  - 19. A plant comprising a cell according to claim 18.

25


- 20. A plant according to claim 19 selected from plants of the families Brassicaceae, Poaceae, Solanaceae, Asteraceae, Malvaceae, Fabaceae, Linaceae, Canabinaceae, Dauaceae and Cucurbitaceae.
- 21. A process for at least partially inactivating a DNA mismatch repair system of a plant cell, comprising transforming or transfecting said plant cell with a DNA molecule according to any one of claims 1-3 or 7-12 and causing said DNA sequence to express said polynucleotide or said polypeptide.
  - 22. A process for at least partially inactivating a DNA mismatch repair system of a plant cell, comprising transforming or transfecting said plant cell with a chimeric gene

according to any one of claims 13-16 and causing said DNA sequence to express said polynucleotide or said polypeptide.

- 23. A process for at least partially inactivating a DNA mismatch repair system of a plant cell, comprising transforming or transfecting said plant cell with a plasmid or vector according to claim 17 and causing said DNA sequence to express said polynucleotide or said polypeptide.
- 24. A process for increasing genetic variation in a plant comprising obtaining a hybrid plant from a first plant and a second plant, or cells thereof, said first and second plants being genetically different; altering the mismatch repair system in said hybrid plant; permitting said hybrid plant to self-fertilise and produce offspring plants; and screening said offspring plants for plants in which homeologous recombination has occurred.
- 25. A process according to claim 24 wherein a first gene is incapacitated in said first plant, a second gene is incapacitated in said second plant, and said first and second genes are incapacitated in said hybrid plant thereby altering the mismatch repair system of said hybrid plant.
  - 25. A process according to claim 25 wherein said incapacitation of the mismatch repair system of said hybrid plant is reversible.
- 26. A process according to claim 24 wherein a new genetic linkage of a desired characteristic trait or of a gene which contributes to a desired characteristic trait is 20 observable in at least one of said offspring plants.
- 27. A process for obtaining a plant having a desired characteristic, comprising altering the mismatch repair system in a plant, cell or plurality of cells of a plant which does not have said desired characteristic, permitting mutations to persist in said cells to produce mutated plant cells, deriving plants from said mutated plant cells, and screening said plants for a plant having said desired characteristic.
- 28. A process according to claim 27 wherein said step of altering the mismatch repair system comprises introducing into said hybrid plant, plant, cell or cells a chimeric gene according to claim 13 and permitting the chimeric gene to express a polynucleotide which is capable of interfering with the expression of a plant polynucleotide sequence in a mismatch repair gene of the hybrid plant, plant, cell or cells, or a polypeptide capable of disrupting the DNA mismatch repair system of the hybrid plant, cell or cells.
  - 29. A process according to claim 28 comprising inactivating an MSH3 gene and/or an MSH6 gene of said plant.
- 30. A process according to claim 28 comprising inactivating an MSH3 gene and an MSH6 gene of said plant.

- 31. A process according to claim 27 comprising at least partially inactivating the mismatch repair system of said plant in a predetermined cell type or in a predetermined tissue of said plant.
- 32. A process according to claim 31 further comprising restoring mismatch repair 5 in said cell type or said tissue.
  - 33. An oligonucleotide capable of hybridising at 45°C under standard PCR conditions to a DNA molecule according to claim 1 with the proviso that said oligonucleotide is other than SEQ ID NO:1 or SEQ ID NO:2.
- 34. An oligonucleotide capable of hybridising at 45°C under standard PCR to conditions to the DNA of SEQ ID NO: 18 with the proviso that said oligonucleotide is other than SEQ ID NO:1 or SEQ ID NO:2.
  - 35. An oligonucleotide capable of hybridising at 45°C under standard PCR conditions to the DNA of SEQ ID NO:30 with the proviso that said oligonucleotide is other than SEQ ID NO:1 or SEQ ID NO:2.

Figure 1






Figure 2

Comments/References: 52= 3' side of S5 (AtMSH3) 1244bp in pUC18/Sma1




Figure 3

Comments/References: 13 = 3' side of S5 (AtMSH3) 2104bp in pUC18/Sma1

4
Ø
ŭ
2
.4
(E)

80	144 15	204 35	264	324 75	384 95	<b>444</b> 115	504 135	564 155	624 175	684 195	744	804	864 255	924
CGCGCGAAAATTGGCAACCCCAAGTTCGCCATAGCCACGACCACGACCTTCCATTTCTCTTAAACGGAGGA	CCC	CCG P	CTC L	GAT	GTT V	AAG K	GGA G	TTC	GCA A	GCA A	GAA E	AAT N	GAA E	GTT
ACGG	GCT A	CCA	CAC H	CCC	TAT Y	CTA L	TTC F	AAT N	AAT N	GGT G	CTT	AGT S	ATT I	GTT
TTAA	TTC	CCG P	GAC D	GTA V	GAA E	GAG E	TTC F	CAC H	GTG V	CAT	ACG T	CAG Q	GGT G	GAA
TCTC	TTC F	ACA	TCC	CCA P	GAG E	GTG V	AGA R	GAT D	$_{\rm L}^{\rm CTG}$	TCC	GCC A	TCA S	TGT	GGT
CATT	CGT R	TCA	CTC	AAC N	CCG P	GTG V	TAC Y	ATG M	AGA R	AAG K	AAA K	GGT G	၁၅၅	ACA
CTTC	TCT	TCA S	CTT L	CAA Q	TCG S	CAA Q	AGG R	CAT H	AGA R	ATT I	ACC T	TTT F	TTA L	TCG
CGAC	ATT I	GAA E	A.A.G K	ACT T	CCC	CAG Q	TAC Y	GCT A	GTG V	GCC	TAT Y	GG'F G	ACA T	A1'T
ACCA	ACG T	GCC	CGT R	CAC H	gaa E	GAA E	GGT G	TAC Y	CAT H	GCA A	$_{ m L}^{ m TTG}$	GAA E	GAG E	GAA
CACG	CAG Q	GTA V	AAG K	CCT P	CTG L	$_{ m L}^{ m TTG}$	GTT V	ATT I	TTC F	ACT T	GCG A	GAA E	TCG S	GTT
TAGC	CAG Q	CCG	TCC S	TCT S	TTT F	CCA	gaa E	GGT G	AAT N	GAA E	TCG S	GGT G	A.A.G K	၁၅၅
CCCA	AAG K	AAT N	CC'I	$_{\rm L}^{\rm CTT}$	AGA R	ACA T	GTG V	TTG L	${f TTG}_{f L}$	ACT T	$_{ m L}^{ m CTG}$	GGT G	GTT V	GTT
GTTC	CAA	CCG P	TCT S	AAA K	CAG Q	TAC Y	ATG M	GTG V	CGA R	CAG Q	GGA G	TGT C	AGA R	GTT
CCAA	AAG K	GAA E	TTC F	CCT	CTC L	AAA K	$_{\rm L}^{\rm TTG}$	CGC R	TTT F	A.A.G K	CGG R	GGT G	GAG E	GGT
CAAC	ე ეეე	CAC H	TCC	AAG K	TTT E	AGG R	GTT V	GCA A	ACA T	GTG V	TTC F	GGT G	GAT D	GTC
TTGG	ATG M	ACT T	GTA V	AAA K	AGA R	TCG S	GTG V	GCA A	CCA P	GTA V	TTT F	AGT S	GTG V	AGA
AAA	AAAGCAATT	CCG P	ACT	CCC	CAA	TCA	GAT D	ATC I	GTG V	GGT G	CCT P	ATA I	GTT V	GTT
ອວອວ	AGCA	TCC S	GCC A	TCA S	CAC H	TCA S	CCA P	gag E	AGT S	ATT I	3 3 6	GAT D	TGT C	GAT
		AAA K	TCC S		TTA L			GCG A	GCG A	AAG K		GAG E	GTT V	TTT
CTAAGAAAG	GATTACGAAT	CCC		GCC A	AAT	gaa E	AAG K	GAC D	ACG	TAC	CGG R	GCT A		AGT
cCTA	GATT	AAA K	AAG K	209		CCC	AGC	GAA	ATG	GGA	AAC	gce A	TTC	ATG
7	81 1	145 16	205 36	265 56	325 76	385 96	445 116	505 136	565 156	625 176	665 196	745 216	805 236	865


984 295	1044 315	1104 335	1164 355	1224 375	1284 395	1344 415	1404 435	1464 455	1524 475	1584 495	1644 515	1704 535	1764 555	1824 575
	<b>,</b> n			£-1	E.	4	£4	ď	ď	O	E	æ	£	ဗ
TTG L	GTG V	AGC S	TTA L	GTT V	CAT H	TCA S	AAT N	GTA V	ATA I	TCC	TTT F	GGA G	GCT A	AGG R
AGC S	${f TTG}_{f L}$	TTC F	AAC	ACA	TGC C	TTG L	A.A.A.	ACA T	TTG L	TCT S	GAG E	CGT R	GAA E	ATG M
TTG L	TTT F	TGT C	GGT	${f TTG}$	TTT F	TCT	GTG V	CTT	AAT N	AGT S	CCT P	CAA	ATG M	GAA E
ATT	AAG K	GAT D	GCA	TGC	ACG T	CGC R	GTT V	ACA T	AGA R	CAT H	TCA S	ATT I	GTT V	rct S
GTG V	GAG E	CTG L	AGC S	TCT S	CTA L	TTT F	GAG E	CAC H	GAT D	TCT	GTA V	GAT D	GCA A	GAC D
GCT A	ACT T	TCA S	ATC I	ATG M	GCC A	TCA S	TTG L	AAT N	TGC	<b>G</b> G <b>A</b> G	ATT I	TCT S	ATT I	CAA O
GAG E	CAA	GCC	AAA K	GGA G	CTC L	GCC	CAG	ATG M	CTA L	ATG M	GCA A	TCA S	TTC F	GGC ATA AAG CAA G I K Q & (Continued)
TTA L	CAA	CGT R	gaa E	AAA K	GCC A	000 0	CAM	AAT N	CCT P	TGC	AGA R	AGA R	GAG E	ATA I Sont
GGA G	TCA	GAA E	TGT C	GAA E	CAA O	CAA	$_{\rm L}^{\rm CTG}$	CAT H	CAT H	GCT A	GAG E	TCT	ACA T	<b>၁)</b> ၁၅၅
AGT	CTT L	GTG V	TTA L	GCT	GTT V	TAC Y	ACT T	TTC F	ACT T	TCT S	TCT	ATG M	GCC A	
AGA R	CCT P	CGA R	TCA S	GCT	ACT T	CTT L	AAT N	TTA L	GTG V	ATT I	GGT G	GCT A	AAA K	cgg ctr R L
ATG M	CAG O	GTT V	ATT I	GAG E	CTG L	ATC 1	GCC A	TCC S	TGG W	GAG E	GAA	ACA	GCT	CAG Q M
${ m TTC}$	၁၅၅	AAC	GTT V	$\mathop{\mathtt{CTG}}_{\mathbf{L}}$	CAT H	AGG R	TCA S	၁၅၅	CAC H	TCT	GAA E	${ m TTG}$	ACT T	ATT I
AAT N	CTT L	TCA S	GAG E	A.A.G K	CCA P	GAA E	$_{ m L}^{ m CTC}$	TCT	AGA R	GTT V	GTT V	GTC V	CGG R	CAA Q
GAT D	$_{ m L}^{ m TTG}$	ACC T	GAT D	ATG M	ATG M	TTT F	ACT T	GAA E	$_{\rm L}^{\rm CTT}$	GCT A	${ m rrg} \ { m L}$	TCA S	CAT H	AAG K
AAT N	CTG L	CCT P	GTA V	gaa E	AAC N	GGA G	ATG M	TCG	$_{\rm L}^{\rm CTT}$	GAT D	GAG E	TCC	TTT F	ອອອ
TTC F	GAG E	GGA G	GCA	AAA K	ATG M	TTT F	GAG E	GGA G	AGG R	CTT L	AGT S	CTC L	ATC I	GCG A
GAG E	GCT A	GCT	AAT N	GAT D	ATT I	CAG O	ACA T	GAT D	TCC S	CGG R	AGC S	GTG V	AGA R	CTT L
GAA E	CCA P	CAT M	GGT G	GAT D	ACA T	AAA K	AAC	TCA S	GCT G	GCT A	CTC L	CTC L	ACA T	TTA L
TAT Y	TCA S	GCA	AAT N	GAA E	CAT H	CTC L	AGT S	AAT N	TAT Y	TCT	CAG Q	TAT Y	ATA I	ATT I
925 276	985 296	1045 316	1105 336	1165 356	1225 376	1285 396	1345 416	1405 436	1465 456	1525 476	1585 496	1645 516	1705 536	1765 556

1884 595	1944 615	2004	2064 655	2124 675	2184 695	2244 715	2304 735	2364 755	2424 775	2484 795	2544 815	2604	2664 855	2724 875
TCA S	GCT A	GCT A	TTT E	TTG L	AAG K	GCA A	AGT S	CAC H	TGT C	CAA Q	ATC I	ATA I	GGT G	GAA
ATT I	GCG A	$_{\rm L}^{\rm CTT}$	TCA S	CAT H	ACC T	CTA L	TTC F	$_{\rm L}^{\rm TTG}$	GAC D	TTA L	ATT I	TCC	GAT D	CTA L
GTT V	GAA E	GAG E	GCT	ACA T	AGC S	GCT A	AGT S	TGT C	GAT D	ATA I	CAA	ATT I	CTT L	TTT F
TCT	AAG K	CCT P	ATA I	ATC I	AAT N	CTA L	A.A.G K	GAC D	GTG V	ACT T	TGC	TTA L	GTG V	ACC
ATT I	AAT N	TTT F	TCG S	9 9	GTA V	GAG E	CTC L	ĊTG L	TTT F	GAG E	TAT Y	GCT	CAC H	AGT
${f TTG}$	CTA L	CAA Q	TCC	TCG S	A A A	GAT D	TTC F	GCA A	GAG E	CTG L	GAA E	GTT V	CTG L	AGA R ed)
AAA K	GCC	GAC D	GAT D	GTG V	GTG V	TTG L	AGT S	GCT A	CCC	GTA V	9 9	CAA O	AAG K	G CAT GGC AG H G R (Continued)
AGA R	TCT S	AGC S	CTG L	CAA O	TGG W		GAT D	CTT L	CGT R	CCT	GAA E	CGT R	GCC A	CAT H
${ t TTG}$	CTC L	TCC S	AAG K	CTT L	AAT N	GCT	TGG W	GCT A	GTC V	CAT H	GCA A	ATC I	TTC	ည်ထ 🐾
$_{\rm L}^{\rm crr}$	CTT L	AC'F T	GAA E	TTT F	ATG H	GTA V	TCG S	CAA Q	TAT Y	cgr R	CAT H	TAT Y	TCA S	AGT ATC S I
ACT T	AAA K	A'I'C I	AGG R	GAA E	CCT P	ATA I	GCT	GTT V	AAC N	GGT G	$_{ m L}^{ m TTG}$	TGC C	GCG A	AGT S Figu
TCT	GGA G	CTA L	A'rc I	$_{\rm L}^{\rm TTG}$	GTC V	GAA E	CGA R	GCC A	AAG K	TCT S	ATT I	AGC S	CCA P	GAC D
CGA R	GCC A	ATA I	GTC V	AAT N	AAG	CCA P	AAC	GCT A	AAC	CAG Q	ACA T	AAG K	GTA V	TCAS
GTG V	AAT N	GAC D	TTA I.	CGA R	TCC	CCC P	GTG V	A.A.G K	AGA R	ATA I	GAC D	GGA G	TTT E	GCT A
ACT	GAC D	$\frac{\text{CTC}}{\text{L}}$	GTT V	ATT I	GAT D	CAT H	ATT I	TTT E	TCT	AAC	AAT N	GGA G	TCC	GGT G
GCA	GTT V	$_{\rm L}^{\rm TTG}$	GCA A	GCT A	GTT V	TAT Y	GCC	GAT D	CTA L	ATA I	CCA P	ATG M	GGT G	ATG
TCT	GTG V	GAC D	CAA	CTC	CCC P	CGA R	CTT L	ACA T	ACT	GAG E	GTC V	AAC	GTT V	CGG R
CAA	GTT V	GGT G	CGC R	AAG K	CTG L	ATT I	CAT H	TAC Y	TCA S	GTT V	TTC F	CCT P	CAG Q	ACT
ATG M	CCT	CGA R	GCT A	AAG K	GAG E	ACT T	GAA E	TAC Y	CTT	CCA P	AAC N	GGA G	GCT A	TTC F
AGT S	TCC S	GTT V	GAA E	CGC R	ATA I	AAG K	ACT T	AGA R	TCC	GAA E	GAT D	ACC	ATG M	GTT V
1825 576	1885 596	1945 616	2005 636	2065 656	2125 676	2185 696	2245 716	2305 736	2365 756	2425 776	2485 796	2545 816	2605 836	2665 856

2784 895	2844 915	2904 935	2964 955	3024 975	3084 995	3144	3204 1035	3264 1055	3324 1075	3397 1082	3458 5	3522 16
GAA GCG TCA CAC ATA ATC AGA ACC TGT TCT TCT TCG CTT GTT ATA TTA 278 E A S H I I R T C S S R S L V I L 895	GGA AGA GGC ACT AGC ACA CAC GGT GTA GCC ATT GCC TAT GCA ACA TTA G R G T S T H D G V A I A Y A T L	CTA GCA GAA AAG AGA TGT TTG GTT CTT TTT GTC ACG CAT TAC CCT GAA ATA L A E K R C L V L F V T H Y P E I	AGT AAC GGA TTC CCA GGT TCT GTT GGG ACA TAC CAT GTC TCG TAT CTG ACA S N G F P G S V G T Y H V S Y L T	GAT AAA GGC AGT TAT GAT CAT GAT GTG ACC TAC CTA TAT AAG CTT GTG D K G S Y D H D D V T Y L Y K L V	TGC AGC AGC TTT GGT TTT AAG GTT GCT CAG CTT GCC CAG ATA CCT CCA	CGT CGA GCC ATT TCA ATG GCT GCA AAA TTG GAA GCT GAG GTA CGT GCA AGA R R A I S M A A K L E A E V R A R	AAT ACA CGC ATG GGA GAA GGA CAT GAN GAA CCG AGA GGC GCA GAA GAA 3204 N T R M G E P E G H E E P R G A E E I 1035	TCG GCT CTA GGT GAC TTG TTT GCA GAC CTG AAAA T'FT GCT CTC TCT GAA GAG GAC 3264 S A L G D L F A D L K F A L S E E D 1055	332 107	ACT TGT TCA TTT TGA TTTAATCTTAACATTATAGCAACTGCAAGGTCTTGATCATCTTGGTTGCG 3397 T C S F *	ATG TGT ATT AGT ATA ACA AGA AAA GAG AAT TAG AGAG ATG GAT TCT AAT CCG M C I S I T R K E N * M D S N P	CAG TAC ATC TTT TCT CCA CCC GCA TAA AAAAAAAAAA
A AGT S	G CTT L	T CTC	G ATC	G AAG K	r crr L	sT ATA I					A( 2)	
TTA AGT L S	GAG CTT E L	CAT CTC H L	ပ္ခ	CAG AAG Q K	GGT CTT G L	TGT ATA C I	AGA AAT R N	ATT TCG I S	TGG AAA W K	CCA ACT P T	TACTAACTT A'	TTG CAG L Q
GAA T E	GAT G	CAG C	GCT GA	TTG	CGT G	TCA 1	GAG P	TCT P	CCT 1	AAA C	TACT	GTG V
2725 876	2785 896	2845 916	2905 936	2965	3025 976	3085 996	3145	3205 1036	3265 1056	3325 1076	3398 1	3459 6

1 MGKQK	2 SDHLAAASPKÄPKLBPHÄQÄPVPDPHLHQÄPLQÄPLRPSPÄRYÄPRTSSBRÄYÄPÄÄQOVPEÄKPDVZÄYPPÄÄÄ 6 VSSKNSKNSKÄTSGTSTÄPÄDIUPAKKLDÄTHKÄKSDENVKAKUDOKKEUTOPÄÄKKANKSPTÄÄLÄPÄDKOVKDUKMURDKZÄVIMÄG	4 POBDÁBIAARVOGITABMOHN 1 EARDAVTVBRITERIKLVPGKLTÍDES	3 filearediscococecopeopeopeopeopeopeopeopeopeopeopeopeopeo	6 ISOQTEKTIVAHAQPTSNVRVERA SLDCYSNGNA VD SVISIOCEKISAGNLEDDK EMRLEAA EKGMSCLTVHTIM NH PELTVQALALTPCHLKOF 8 IPPLHVA KPERDISCRIIHKOLYDLEDHVVQAIRVANIKIONSPELTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTT	1 PERILY COASTRELS SMTRHTUSART COOPS VVK NN SDGS ES GSTRHNAN HOLLV KOSRLER EN EVOT SPECORAL ES AR HOLDV SKINNS B NOOVAL IF SITS FOR	6 QLSSELVEROSKRAIVSPRYLVLSSVLTAMSKSSDIQBUITETFHRUAKATSVIAVMERULAGKQIGRLOIKQDSEMRSMQS-ATVKSTTUR 7	1 President Structures and Contraction of the Contr	30 VDSKVPMNÄVKVRSTRYBPPETVAGLDE GALATERGAIVNNASMOSFIKSPBRYTTOPTBAVGA GALAGGG, MSESTLSARKKYVRPTPVD 36 Qikdipoden KVRMRMV SRTTTPRTOKLTOR BYYKDLIIHESELOYKE FINKITAR YTTEN DAGY OGILSIAATSCENVRYRRAT PVM G	75 CEPVRÍNIQSORHPVLETICODNY VPNDTICHABGEYCQIITGPNHGGKSCYIRQVALICHHAQYGSPVPABYARLICHTRIGAHDOCHTRAGASDSCOM 83 QOAliakharapapiies-Lovyyvendinmspenckiniitgpnhggkskyirovalithaqigspvpabyirboliterigahdding	70 RSTELEELSRASHIIRTOSSRSIVILDELGROTSTEEDGVAIAYAALLAEKRO-BYGEVIRIPPRIADIKSPLIRMYHEDKVEKOKTOK 73 OSTEKVERLDILHEIKROMARSELLEDEVOROTOTABRAISYALIKYVSELSDOPELEFITTIPPALODIKSPLIRMYHEDKVEKOKTOK	64 Suddyt isykbyrgicsrsygeryngippschrismarfraðvharrhyderepegarssergerserser 63 Sahsút flykbyrgigarsygneringiskeideretirismarfrasin	S9 APEFLEHAMEHAGET BEF
##	9.00	134	213	9 9	4 2 8	696	5.0 5.0 1.0	99	77 5	T 89	99	105
E E E E E E E E E E E E E E E E E E E	SH3_A	S 23 3	S H 3	8 8 E E	HSH3. ACHSE3. SC	E SE D	ESB3_AC	S H 3	SE 3. A. S.	SH3.A	S E 3 - A	K SE 3 AC K SE 3 SC

Figure 6






Figure 7

Comments/References: 43= 5' side of S8 (AtMSH6) 2182 bp in pUC18/Sma1




Figure 8

Comments/References: 62= 3' side of S8 (AtMSH6) 1379bp in pUC18/Sma1

(I)
н
-
_
ס
2.
21

80	153 4	213 24	273 44	333 64	393 84	453 104	513 124	573 144	633 164	693 184	753 204	813 224	873 244	933 264
၁၁၁၁	ניז	TCC	999 999	aga R	CCG P	AAA K	GAT D	AAA K	gaa E	GCT A	GTA V	GAA E	AGA R	TTC
rtcc	c cAG	GTT	GAA	GTT V	AAG K	GTA V	AAT N	GGT G	GTA V	CGT R	CCT P	AAA K	AAT N	GTT V
ATCT	G CGC R	TTG	AAG R	GAG E	TTT F	TTT F	CTG L	AAT N	TCA S	CCA P	GTT V	AAG K	GCC	GAT D
GCCA	G CAG	GGT	GTG V	GAT D	GGA G	AAG K	CCG P	AAT N	AGA R	CGI R	AAG K	GAG E	GAT D	CCT P
aaaagttgagccctgaggagtatcgttttcgccatttctacgacgcaaggccaaatttttggcgccaatcttccccc	TTTCGAATTCTCTCAGGTCAAACATCGTTTCTCTCTCTCT	AAG K	A.A.T N	GTC V	TCT	CAT H	GTT V	TCC	CTT L	ATG M	GAT D	GGA G	AGG R	CCA
TTT	AAAA	ACG	TTT F	TCT S	CCG P	ATG M	GTT V	CGT R	GAA E	999 9	GAG E	TGT C	ATC	ATA
AAAA	TTCC	ACT	CGA R	AAA K	CTG L	ATT I	GAT D	T'I'T F	GCT A	CCA P	AAG K	GTT V	CGA R	CAC
ງວງງ <u>າ</u>	ACAA	GCG A	CCA P	TCG S	GTC V	AAT N	GAN E	CAA	AGA R	ACA T	TTT F	CCG P	TCT	TTA L
GCAA	TCTC	GCG A	GGA G	GTT V	CGT R	TCC	CGA R	CCT P	ე ე	gaa E	ACT T	GAT D	TCT	ACC T
CGAC	TCIC	ACC T	GGA G	GCT A	CGT R	TTC E	AGC	ATT I	AGT S	CCA P	ATG M	CAG Q	GAG	, AAG K
TCTA	TCAC	CCC	AGC S	TTT F	CCG P	$_{\rm L}^{\rm cTG}$	AGG R	GT'r V	TTC F	GGT G	GAA E	CTC L	CTT	AGA R
CATT	TCTC	AAA K	ე ეეე	CGT R	GTT V	TCC S	GAG E	GAT D	AGT S	CCT P	GAT D	ATG	TGG W	GAT D
ວອວວ	TTTC	CAA	၅ ၁၅၅	GTA V	AAG K	TCG S	GGA G	AAT N	TTT F	GTT V	GAG E	AAA K	GAA	. TAC Y
GTTT	ATCG	TTC	9 999	TCT	GAG E	GCT A	TCT S	GCT A	GCT A	GAT D	CTG L	CTG	TTT F	CTT L
TATC	AAAC	TTC	၁၅၅	GCT A	CCG P	GAT D	TGT C	AAG K	CAT H	3 3 9	GTT	AGG R	AAA K	- CCC
GGAG	CTCP	TCT	AGC S	GAC D	CCA P	GST G	GAT D	ATG M	AAC N	GAT D	CGA R	AAA K	ACC T	GAT
CTGP	TCAG	$\mathop{\mathrm{TTG}}_{\mathbf{L}}$	GCT	၁၅၅	ACT T	GCC	CGA R	TGT C	AGA R	GTA V	AAG K	AAC	GGA	GAT D
:AGCC	TCTC	ATT I	GCT A	A A A	GAT D	TCC	GAT D	$c_{\mathbf{L}}$	GAA E	GGA G	TTG	TCT		CCT
GTTG	GAAT	TCG S	GAT D	GCT A	ACG	GAA E	GAT D	TCT S	CAA O	ATA I	CGC R	GAC	AAC	. CGT
AAAA	TTTC	AGA R	၁၅၅	GAT	GGA	GCT A	GTC	TCA	ACT	GAT D	TCT	$_{\rm L}^{\rm TG}$	GTA V	AGA R
-	81	154	214	274	334 65	394 85	454 105	514 125	574 145	634 165	694 185	754	814	874 245

993 284	r 1053 304	1113	1173	4 1233 364	3 1293 384	4 1353 404	2 1413	4 1473 444	3 1533 464	r 1593 484	л 1653 504	5 1713 524	T 1773 544	G 1833 564
ATT I	GGT G	GGT	GTT V	ATA I	9 999	TCA	ATC	GAA E	ACG T	GAT	TCA	GAG E	TTT F	GAG
GAC I	TTA L	GTT V	AAA K	ACT T	ATC I	TGT C	TCC S	AAG K	TAT Y	ACA T	GAA E	GGA G	ATT I	CTT
ATG G	GAA E	CAG Q	TAT Y	AAT N	AAC	AAG K		CCA P	aaa K	GAT D	TCT S	CTT L	GAT D	AAT N
TAT A Y	GCG	AGA R	GGA G	GCT A	GGA G	CAA	GTT V	TCT S	AGG R	999 9	TCT S	GCT A	9 999	GTA V
GAA T E	GAT D	TGC	CGT R	GGT G	GAG E	CTA L	TGG W	GTT V	CTA L	ATG M	GGT G	AGT S	CAT H	ATG M
AGT G S	CTA	AAA K	GCT A	AGA R	AGC S	GAG E	TTT F	CAG Q	GCT A	GTA V	A.A.A	CTT L	A.A.G K	ACG
AAG A K	GAG E	GGA G	TTA L	GCC A	GCA A	ATG M	AGG R	ATG M	A.A.G K	CAA O	TTT F	GCC	CTT L	CAG 0
GTT A V	TAT Y	GTG V	CTA L	AAA K	ACA T	AAA K	${ m TTG}$	${ m TTG}$	CAA O	CCA P	TAC	GTT V	GTA V	၁၅၅
AGT G S	$\mathop{\tt CTG}_{\rm L}$	GGT G	aag K	GCA A	TCA S	ATC I	GCC	TTA L	GCA A	GTA V	GGA G	GAT D	GAT D	GAT D
TGG A W	GAG E	AGT S	CAA Q	CAA Q	CCA P	GAG E	GCT A	GCG A	GAA E	CCA P	AAC N	TGT C	GAA E	ATT I
TAT T Y	TAT Y	ATG M	GTG V	GAC D	ACT T	AAA K	TGT C	GGA G	AGA R	GCT A	TCT S	GAA E	CTA L	AGA ATT R
CAA T	TTT F	ACC T	GCA A	TCT S	TTA L	ATA I	GAC D	CTT L	TCA S	TTG L	GAA E	AAT N	A.A.G K	CTC 1.
AAG C	AAA K	ATG M	GAG E	ACA T	GTA V	GCT A	GTT V	GCT A	CTA L	CAG Q	ATA I	CTA L	CTA L	TGT
CAA A	999 9	AAG K	GAT D	GAA E	CAG Q	$_{\rm L}^{\rm CTT}$	TTT F	GCT A	9 9	GTA V	ATA I	GGT G	AGG R	GGT
rca c	GTG V	TGG W	ATA I	CTA L	GTT V	$_{\rm L}^{\rm CTT}$	GCT A	TGT C	AAA K	GCG A	AAT N	GAT D	TCT S	AGG
SCA T	AAA K	GAC	999 9	CAG Q	CTA L	CAT H	TTT F	TCA S	AGT. S	ACG T	AGA R	GTT V	CTG L	TAC Y
rct G s	TTT E	CTT L	AGT S	GAG E	AAG K	GTC V	GGA G	GCA A	GAC D	TCT	GTT V	GCT A	CAT H	GTT V
-	TTC F	GAG E	GAA E	ATC I	AGG R	GCC	TAT Y	GAT D	TAT Y	999 9	GGA G	TGT C	AAT N	CAA
AA A K	CTT	AAG K	TCT	CGA R	CCA P	GAT D	GTG V	GAT D	TTA L	ACA	GCT A	AAC	ATT I	TAC
AAG AAA ATG K K M	GTG (	CAC	ATC	GGA	ATT	CCT	ACT	AGC	GTG	TTG	GCT	TGG	CTA	CCA
934 265	996 285	1054 305	1114	1174 345	1234 365	1294 385	1354	1414	1474	1534 465	1594 485	1654 505	1714 525	1774

1893 584	1953 604	2013 624	2073 644	2133 664	2193 684	2253 704	2313 724	2373 744	2433 764	2493 784	2553 804	2613 824	2673 844	2733 864
AAC	GAT D	AGT S	CGC	GTG V	GAT D	AAA K	GCC A	ACT T	CAC H	GGA G	ACA T	ggr G	ATT I	cgr R
GAT A	AAA G K	GAA P	GGA C	AAA K	ATT ( I	TGT 1	GCA A	GAA	ATT I	GCT A	AAA K	GAT	AGC S	CTT L
CTT	CTC L	TCA S	CTC	A A A	GGA G	CTC L	GAA E	GCT A	GTC V	TCT	CAG Q	GCC	၁၅၅	CTT L
TAT Y	CCA P	AAC N	CTG L	999 9	AGT S	AAA K	TTC F	AAC	GAG E	CTC	AAT	GCA A	AGT S	ACT T
A.A.A	CAT H	GCA A	AGA R	CTG L	AGA R	TAT Y	CAA Q	GAA E	TCT	AGT	cAG	GTT V	A AGC S	A TCA S
TAC Y	TGC C	ACG T	GAA E	CTT	1.TC F	CTT L	TCT	. GAT D	TGG W	GCA A	GAT D	GCA	A AGA R	A AAA K K
TTG L	ATC I	TTC F	T'f A L	GCT A	999 9	TTG L	CTT L	ACA T	CAA	GCA A	ACA	TTT	. AGA R	GGA G inued
ACC T	TGG W	GAA E	GAC D	CC.T P	X X	AGT S	TTT F	GTG V	ACT T	ATC I	GCT A	CCA P	GCT A	G GGC GGA G G G (Continued)
999	A.A.T. N	gaa E	CCA P	TTG L	GTG V	ATG M	TTA L	GAT D	GCA A	GCA A	GAA E	CAT H	GAG E	M M
TCA S	AGG R	GTT	CTT L	GTG V	ATT I	A1'G M	GAG E	CAA	AGA R	TTT F	TCA S	TGG W	၁၅၅	AAC N gure
CCT P	TTA L	GTA V	A.A.A K	TCT S	CAA	AAT N	CTA L	AAC	GAA E	TCT S	GAA E	CTA	CTT	CCA P Fig
GGT G	CTC L	GAT D	CAC	GCC	000 0	TCA S	999 999	CAG Q	ATC I	AGA R	CCC	GGA G	CTC	GGA G
GGT G	CGA R	CTT L	CTC 1.	TCA S	TTT F	GAA E	AGC S	TAT Y	TTT E	CTG L	TTT F	CAA	ATA I	ACG T
GAT D	A.A.G K	CGG R	TAT Y	TCA S	GCA A	AAG K	AAA K	AAT N	CTT L	GTC V	ATT	ATC I	GAT	CTG L
TGT C	GGT G	A.A.A K	CAG Q	CGA R	AAA K	CAG Q	GGA G	CCA P	GAA E	GAT D	GTT V	AAA K	AAT N	TTA
AGC S	ACT T	AAT N	၁၅၁	GTT V	GTT V	CTA L	GTA V	TTT F	ATC I	CTA L	CCT P	CTT	P CCG	, TTG
AAT N	CCA P	ATC I	ACT T	AGC S	CGA R	GCT A	TTA L	GAC D	CTT L	TGC C	AGG R	ATA I	GTT V	S
AAC	AGT	AGC S	ATC	TCT S	CAA Q	${ m TTG}_{ m L}$	ATA I	AGC S	ATA I	AGC S	9 8	CCA P	CCT	000 8
TTT	GTT V	GAA		AAG K	A.A.A K	$_{\rm L}^{\rm TTG}$	CCT P	GAT D	ACA	ATA I	ATG M	999 999	TTG L	CCT P
ATA	TGT	GTA V	ATG M	ATC I	CTG L	CTG L	CTT	ATA I	CTC	ACC	AGC	AAA K	CAA	CAT H
1834 565	1894 585	1954 605	2014 625	2074 645	2134 665	2194 685	2254 705	2314	2374	2434 765	2494 785	2554 805	2614 825	2674 845

2793 884	2853 904	2913 924	2973 944	3033 964	3093 984	3153 1004	3213 1024	3273 1044	3333 1064	3393 1084	3453 1104	3521 5	3579 19	3606 28
TAC GTG CCG TGT GAG TCT TGC 27	GCA TCT GAT AGA ATC ATG ACA A S D R I M T	TCA GTT CTT CAG AAT GCA ACT S V L Q N A T	ACT AGT ACT TTC GAT GGA TAC T S T F D G Y	GTT CAA TGT CGG ATG CTC TTT V Q C R M L F	CAC CCA CGT GTC ACC TCG AAA H P R V T S K	CCA CGT GGT TGT GAT CAA GAC P R G C D Q D	GAG AGC TAC GGA CTT CAA GTG E S Y G L Q V	GCA TCA GGT GCT GCA GCC A S G A A Q A	CTA AGA TCT GAG TTC TCA AGT L R S E F S S	CGA GTC GCC CAC AAC AAT GCC R V A H N N A	TGG CAT GAG ATC AAA TCC TCT W H E I K S S	CACTATCTGAAGCTCGTTAAGTCTTTTGCCTCTCT 3	AAA TTG TTT CCT CGA TTA AAA K L F P R L K	3 (Continued)
C TGC	T GGC	ACA GCG T A	AGA GGA R G	GAG AAA E K	GCG TCT A S	TAT CAA Y O	TGT CCT C P	GAA ACA E T	AGT GAG S E	ATT TCT I S	TGC TTA C L	ACTAT	AAA K	g (Con
CTT GGC L G	AGG CTT R L	GAG AC	GGC AC	GTA GA V E	TTC G( F A	GAT T	GCT TO	GT'F G	TCA A	GGT A	TTT T F C	TAA	TAT ATC Y	figure 9
CAA C	ACA P	ACT G	CTG (	CTG (	GAA E	TCT	GGA G	GTG V	AAG K	GTG V	$\mathop{\mathrm{TTG}}_{L}$	ACA T	TTA T	_
000 <b>9</b>	TTC	TGC	GAA E	CAC H	AAG	AGA R	GAG E	CAA	TTC	A TTG	C ACT	r ATG M	TGC T	A AAA K
TTT F	ATC I	GAA E	GAC D	. CGT	ACC T	A TCA S	A ACC T	A AAC N	A AAC N	s TCA S	c GAC D	G GCT A	AAA '	aaa aaa K K
ATC I	ACT T	GTA	CTT	TTT F	r CTC L	A AAA	r TTA L	A CCA	G GGA E	C AAG K	c TAC Y	A ATG M	AAA K	AAA AA K K
GTT V	GAT D	${ m TTG}$	ATC	GTT	CCT P	TTC F	CGT R	A ATA I	r 666	3 CTC L	T GAC D	A TAA	CTT	AAA AA K K
GCC	GTG V	TTT F	GTA V	TCG	CAC	GCA A	TAC Y	GGA G	A A'I''F I	TGG W	A GAT D	AAAA K	CCT	
CTG L	CTC	ACC	CTA L	TAC	TAC Y	TGC	TTG L	GCT A	r TCA	A GAC D	GAA E	r ccc	Ţ	A AAA K
TGT	TCC	AGT S	TCA	GCA A	CAT H	GCT A	TTC	ATG M	: AGA R		ე <u>ე</u> ნ ე		TTT AS	A AAA K
ACA	ATC				ACA	ATG M	GTG V	CTC	AAG K		; ATT I	: TGT C	ATG 1	AAA AAA ! K K !
GCA A	GAA E	GGA	CAG	000 8	GCA	CAC	CTA	GCA	ATG	crg	CCC	TAC	ر ن	A A
2734 865	2794 885	2854 905	2914 925	2974 945	3034 965	3094 985	3154 1005	3214 1025	3274	3334 1065	3394 1085	3454	3522	3580

ACMBRS-RA. 1	almens – at — 69 Prekveravlesofferenden Bulken. – - fvkvotud. –	ACHSES-mi 163 VEDIQVDQDVFQPETFQHEGNIEDENTFREUKVZVLUSZKRIKHLUDVVCQEGYEVERHUSSERIGGA 1700 VEDILLIDER SPETFILDER SPEVFEG Kobé, yeart 210 lakttskkysymteusssöytrmismonskkys frudalskyr <u>u</u> yshygyskysskyskyrgmenyometyskkysymteusssöytrmismonskysk frudalskymmus	aemske-mi 260 saso <u>kovn</u> sv <u>kstampivletky propicileden kelomkatus</u> svero vojisky ouezar svente spoakan. Damini proleduju mske "yoset 112 trrkovnik skampilivletko senjakovil senjakovil sulika sudnamu olasiju sa senjam kunda kunda kunda	5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5	ACHSHS-mi 462 KYTLTOSTAVGLAPVPQVMODT DAAOVRMINEGROOFINGAVDULMEC DIJALGNIJR NEWENSERNINGAPVROOFINGARIOOFINGAN Neme yeert 522 pmaapmaitmevrauerfy Dobrtyae Higheris volkay votor kijopinjy Bo Byyying mandamany mandaman Lighing.	Atmans-mi sea <u>Huberpongeprongeproncvept Dira</u> linmälichpykoværmaradovjerftambermitengeringenerandengeneranden ser mens yeart era Hiberponsonorden konden ser om ser men ser om se	AKMEME-B. 648SVRSS	atmans—mi 725 idadepm	almans.mi eoz sktrupiensoodminspavnadoolevpribiildosamussusiihpnaletagasteraatskuskoodsesteraansestaksendada mans, yeere sis olauponeralminspanalungeraksteraaksesteraansesteraassaatskuskoodsesteraansesteraansesteraanse	ALMSHG-MI 905 SESTELVECTELASVEONATODEEVILDEEGAGTETPOTATASVERILVSKYOCKHIEATHÄMPETKETASKERASKERASVERASVERALLUSKY Meng yerel 1016 Sastelveraskaldunatakassassassassassassassassassassassassas	atment-mi 1010 mitogacembyteromentaunvertaunvoamkgalombykseschmentmelymiskoo-rhmerentme	
--------------	---	--	---	---------------------------------------	--	---	--------------------	-----------------------	---	---	---	--

TTTTTTGGTTGCTAACAATAAAGGTATACGGTTTTATGTCATCAATATAA	50
CTATATATAAAAGAAATGAAAGATATATTATTTTTTTTTT	100
AAAACAACAAGACTTTTTTTTTTTTTTTTTTTTTTTTTT	150
GATAAACGACATCGTTTAATCATTTCCCAATTTTACCCCTAAGTTTAACA	200
CCTAGAACCTTCTCCATCTTCGCAAGCACAGCCTGATTAGGAACAGCTTT	250
ACCATTCTCATATTCCTGAACTACCTGAGTCCTCTCATTGATCTGTTTCG	300
CCAAATCCGCTTGTGACATCTTCTTCTCCAATCTCGCTTTCTGTATCATC	350
AACCTCACCTCTGCTTTCACACGATCCATCGCCGCAGGCTCTGTTTCTTC	400
TTCCAGCTTCTTCGTGTTAATCACCGGAACCGCCGTAGATTTCCCCTTTT	450
TGTTCGAACCGCCATCGAATTTCTTAACCGTTTGAACCGCGACACCGTTT	500
CTCAGAGCTGCGTTAACCGCTTTCGGATCGCGTAGGTCTTGGCTCTTTTG	550
TTTTGATTTGTGGAGAACTACTGGTTCCCAGTCTTGTGTTACTGCTCCTG	600
GGTATCTGCTCGGCATCGTCGATGAATTGAGAGAAAGGAACAACGCGAAA	650
ATTTTATTAATCTGAGTTTTGAAATTGAGAAACGATGAAGATGAAGAATG	700
$\tt TTGTTGAGAGGATTGTGATATTTATATATACGAAGATTGGTTTCTGGAGA$	750
ATTCGATCATCTTTTCTCCATTTTCGTCTCTGGAACGTTCTTAGAGATG	800
ATTGACGACGTGTCATTATCTGATTTGCAGTTAACCAATGCTTTTTGGGT	850
TGGATTCGTGGTACACCATATTATCCGATTTGGCTCAATGGTTTTATATA	900
AATTTGGTTTTCGGTTCGGTTATGAGTTATCATTAAAATTAAGCTAACCA	950
AAAATTTTCGTAAAATTTATTTCGGTTTCAATTCGGATCCCTTACTTCCA	1000
GAACCGAATTATTCGAAACCGGGGTTAGCCGAACCGAATACCAATGCCTG	1050
ATTGACTCGTTGGCTAGAAAGATCCAACGGTATACAATAATAGAACATAA	1100
ATCGGACGGTCATCAAAGCCTCAAAGAGTGAACAGTCAACAAAAAAAGTT	1150
${\tt GAGCCCTGAGGAGTATCGTTTCCGCCATTTCTACGACGCGAAGGCGAAAAT}$	1200
TTTTGGCGCCAATCTTTCCCCCCCTTTCGAATTCTCTCAGCTCAAAACATC	1250
GTTTCTCTCACTCTCTCACAATTCCAAAAAATGCAGCGCCAGAGAT	1300
CGATTTTGTCTTTCCAAAAACCCACGGCGGCGACTACGAAGGGTTTG	1350
GTTTCCGGCGATGCTGCTAGCGGGGGGGGGGGGGGGGGG	1400
${\tt AATGTGAAGGAAGGGGATGCTAAAGGCGACGCTTCTGTACGTTTTGCTGT}$	1450
${\tt TTCGAAATCTGTCGATGAGGTTAGAGGAACGGATACTCCACCGGAGAAGG}$	1500
TTCCGCGTCGTGTCCTGCCGTCTGGATTTAAGCCGGCTGAATCCGCCGGT	1550
${\tt GATGCTTCGTCCCTGTTCTCCAATATTATGCATAAGTTTGTAAAAGTCGA}$	1600
${\tt TGATCGAGATTGTTCTGGAGAGAGGTACTAATCTTCGATTCTCTTAATTT}$	1650
${\tt TGTTATCTTTAGCTGGAAGAAGAAGATTCGTGTAATTTGTTGTATTCGTT}$	1700
GGAGAGATTCTGATTACTGCATTGGATCGTTGTTTACAAATTTTCAGGAG	1750
${\tt CCGAGAAGATGTTCCGCTGAATGATCATCTCTATGTATGAAGGCTA}$	1800
ATGATGTTATTCCTCAATTTCGTTCCAATAATGGTAAAACTCAAGAAAGA	1850
AACCATGCTTTTAGTTTCAGTGGGAGAGCTGAACTTAGATCAGTAGAAGA	1900
TATAGGAGTAGATGGCGATGTTCCTGGTCCAGAAACACCAGGGATGCGTC	1950
CACGTGCTTCTCGCTTGAAGCGAGTTCTGGAGGATGAAATGACTTTTAAG	2000
GAGGATAAGGTTCCTGTATTGGACTCTAACAAAAGGCTGAAAATGCTCCA	2050
GGATCCGGTTTGTGGAGAGAAGAAGAAGTAAACGAAGGAACCAAATTTG	2100
AATGGCTTGAGTCTTCTCGAATCAGGGATGCCAATAGAAGACGTCCTGAT	2150
GATCCCCTTTACGATAGAAAGACCTTACACATACCACCTGATGTTTTCAA	2200

Figure 11

	2250
GAAAATGTCTGCATCACAAAAGCAATATTGGAGTGTTAAGAGTGAATATA	2300
TGGACATTGTGCTTTTCTTTAAAGTGGTTAGTAACTATTAACTATTACGTTATG	2350
CAATCCATTTCCTCAATGTGATTTGTTCACTTACATCTGTTTACGTTATG	2400
CTCTTCTCAGGGGAAATTTTATGAGCTGTATGAGCTAGATGCGGAATTAG	2450
GTCACAAGGAGCTTGACTGGAAGATGACCATGAGTGGTGTGGGAAAATGC	2500
AGACAGGTAAATTAGTTGAAACAACTGGCCTGCTTGAATTATTGTGTCTA	2550
TAAATTTTGACACCACCTTTTGTTTCAGGTTGGTATCTCTGAAAGTGGGA	2600
TAGATGAGGCAGTGCAAAAGCTATTAGCTCGTGGGTAAGGGAACCATCAT	2650
ACTTTATGGAATTCGTTTACTGCTACTTCGGCTAGGATTTAAGAAATGGA	2700
AATCACTTCAAGCATCATTAGTTAGGATCCTGAGAACTCAGGATGTTTTC	2750
TTATTCGTTATATAATAAGTCTTTTCATCAAGGAGTAACAAACA	2800
GCACAATATTTGTGTGCTCACTGGCAAGGCATATATACCCAGCTAACCTT	
TGCTAGTTCACTGTAGTAACAGTTACGGATAATATATGTTTACTTGTATG	2850
TGGTACCCTCATTTTGTCTCTCATGGAGGCTTTGAAGCCTTGTGTTGAAA	2900
CTGGATAGTTACATATGCTTCCAACAGAAACTAGCATGCAGATTCATATG	2950
CTTTCCTATTCTACTAATTATGTATTGACACACTCGTTGTTTCTTTTGAA	3000
AGATATAAAGTTGGACGAATCGAGCAGCTAGAAACATCTGACCAAGCAAA	3050
AGCCAGAGGTGCTAATACTGTAAGTTTTCTTGGATAGGTCAAGGAGAGTG	3100
TTGCAGACTGTTTTTGATCATTTCTTTTTTTTGTACATTACTTTCATGCTG	3150
TAATTAACTCAATGGCTATTCTGGTCTGATTATCAGATAATTCCAAGGAA	3200
GCTAGTTCAGGTATTAACTCCATCAACAGCAAGCGAGGGAAACATCGGGC	3250
CTGATGCCGTCCATCTTCTTGCTATAAAAGAGGTTTGTTATTTACTTATT	3300
TATCTTATCATGTTCAGTTCATCCAAGTCCTGAAAAATTACACTCTTCTT	3350
TACCAATCTTCCATCAAGCTGTGTAAAGGATTTGGAATTAGAAAATCATT	3400
ATTTGATGCTTTGTTTTATATGCAAGAGGTTCCCTTGAAAAGATCTGTTT	3450
AAGATTCTTTGCACTTGAAAAATTCAATCTTTTTAAGTGAATCCCCTACT	3500
TTCTTACAATGATCATAGTCTGCAATTGCATGTCAAGTAATATCATTCCT	3550
TGTTACTGCATCCCCCTCTTTCTTAATGACCATTGTCTATGTTGTGTTTG	3600
TCTCGTGTGCTGGAGAAAATGATAGCTGATCCAAGCTGTACATTATCATG	3650
ATTAAGTAGCTGCTCAGGAATTGCCTTTGGTTACATTGCCTAATGGTTTG	3700
ATGTCAATTTTCTTCTGAATCTTTATTTTAGATCAAAATGGAGCTACAA	3750
AAGTGTTCAACTGTGTATGGATTTGCTTTTGTTGACTGTGCTGCCTTGAG	3800
GTTTTGGGTTGGGTCCATCAGCGATGATGCATCATGTGCTGCTCTTGGAG	3850
CGTTATTGATGCAGGTAAGCAAGTGTATTCTGTATCTTATGTGTACCATG	3900
TGACTTCCTGTGCATATATTTGGGTTGCAGGAACTAATTCTGAATCACCA	3950
TTTGGTATGTTTTTCCAGGTTTCTCCAAAGGAAGTGTTATATGACAGTA	4000
AAGGTAAACTGCTTGTATCGCCAGTTGTTTTGTTAAACAGAATTTAAGGT	4050
AAATGACACTGGTTAATTTAAAGTGCATACATGTTGAAATATTGCAGGGC	4100
TATCAAGAGCACCAAAAGGCTCTAAGGAAATATACGTTGACAGGTACC	4150
TATCAAGAGAAGCACAAAAGGCICIAAGGAAATATACGTCBCCGAATGATAGG	4200
ATTTCAGTAGGCAAGCTAACTGACAATTTAACCGCTCACCGAATGATAGG	4250
TCTCTTAAACATTGCTAATGTAGATGATGTTTATGTTTCAATCTAATAGG	4300
GTCTACGGCGGTACAGTTGGCTCCAGTACCACAAGTAATGGGGGATACAG	4350
ATGCTGCTGGAGTTAGAAATATAATAGAATCTAACGGATACTTTAAAGGT TCTTCTGAATCATGGAACTGTGCTGTTGATGGTCTAAATGAATG	4400
TCTTCTGAATCATGGAACTGTGCTGTTGATGGTCTAAATGAATCTGTGTGTG	


Figure 11 (Continued)

CASCATA A TEATETETET AGGETAAAGG	4450
TGCCCTTAGTGCTCTTGGAGAGCTAATTAATCATCTGTCTAGGCTAAAGG	4500
TGTGTTGGCTTGTTTAGTTTTTGCTTTTCACAAATTAAGCAAAGGAACTT	4550
TTCATAACTTACAGTTTCTATCTACTTGCAGCTAGAAGATGTACTTAAGC	4600
ATGGGGATATTTTTCCATACCAAGTTTACAGGGGTTGTCTCAGAATTGAT	4650
GGCCAGACGATGGTAAATCTTGAGATATTTAACAATAGCTGTGATGGTGG	4700
TCCTTCAGGCAAGTGCATATTTCTTTTTTGATAACTTCAACTAGAGGGCA	4750
GACATAGAAGGAAAAATTCTAATACTTCGTACGGATCTCCAGTAAGTA	4800
AGCCGATTTTGTTTACCTATGTAGGGACCTTGTACAAATATCTTGATAA	4850
CTGTGTTAGTCCAACTGGTAAGCGACTCTTAAGGAATTGGATCTGCCATC	4900
CACTCAAAGATGTAGAAAGCATCAATAAACGGCTTGATGTAGTTGAAGAA	4950
TTCACGGCAAACTCAGAAAGTATGCAAATCACTGGCCAGTATCTCCACAA	5000
ACTTCCAGACTTAGAAAGACTGCTCGGACGCATCAAGTCTAGCGTTCGAT	5050
CATCAGCCTCTGTGTTGCCTGCTCTTCTGGGGAAAAAAGTGCTGAAACAA	
CGAGTAAGTATCAATCACAAGTTTTCTGAGTAATGCCTTCCATGAGTAGT	5100
ATAGGACTAAAACATTACGGGTCTAGCTAAAGACTGTTCTCCTTCTTTTG	5150
CAATGTCTGGTTATTCATTACATTTCTCTTAACTTATTGCATTGCAGGTT	5200
AAAGCATTTGGGCAAATTGTGAAAGGGTTCAGAAGTGGAATTGATCTGTT	5250
GTTGGCTCTACAGAAGGAATCAAATATGATGAGTTTGCTTTATAAACTCT	5300
GTAAACTTCCTATATTAGTAGGAAAAAGCGGGCTAGAGTTATTTCTTTC	5350
CAATTCGAAGCAGCCATAGATAGCGACTTTCCAAATTATCAGGTGCCCAT	5400
CTATCTTTCATACTTTACAACAAAATGTCTGTCACTACTCAAAGCAATGC	5450
ATATGGCTTAGATCTCAACTCACACCCCGAGGATCCTAAAGGGATTTGCT	5500
TTTTATTCCTAATGTTTTTGGATGGTTTGATTTATTTCTAACTTGAACTT	5550
ATTAATCTTGTACCAGAACCAAGATGTGACAGATGAAAACGCTGAAACTC	5600
TCACAATACTTATCGAACTTTTTATCGAAAGAGCAACTCAATGGTCTGAG	5650
GTCATTCACACCATAAGCTGCCTAGATGTCCTGAGATCTTTTGCAATCGC	5700
AGCAAGTCTCTCTGCTGGAAGCATGGCCAGGCCTGTTATTTTTCCCGAAT	5750
CAGAAGCTACAGATCAGAATCAGAAAACAAAAGGGCCAATACTTAAAATC	5800
CAAGGACTATGGCATCCATTTGCAGTTGCAGCCGATGGTCAATTGCCTGT	5850
TCCGAATGATATACTCCTTGGCGAGGCTAGAAGAAGCAGTGGCAGCATTC	5900
ATCCTCGGTCATTGTTACTGACGGGACCAAACATGGGCGGAAAATCAACT	5950
CTTCTTCGTGCAACATGTCTGGCCGTTATCTTTGCCCAAGTTTGTATACT	6000
CGTTAGATAATTACTCTATTCTTTGCAATCAGTTCTTCAACATGAATAAT	6050
AAATTCTGTTTTCTGTCTGCAGCTTGGCTGCTACGTGCCGTGTGAGTCTT	6100
GCGAAATCTCCCTCGTGGATACTATCTTCACAAGGCTTGGCGCATCTGAT	6150
AGAATCATGACAGGAGAGAGTAAGTTTTGTTCTCAAAATACCAATTCCTC	6200
GAACTATTTACTCAGATTTTGTCTGATTGGACAAGGTGGTTTTGCTTTTT	6250
TTTAGGTACCTTTTTGGTAGAATGCACTGAGACAGCGTCAGTTCTTCAGA	6300
ATGCAACTCAGGATTCACTAGTAATCCTTGACGAACTGGGCAGAGGAACT	6350
AGTACTTCGATGGATACGCCATTGCATACTCGGTAACCTGCTCTTCTCC	6400
AGTACTTTCGATGGATACGCCATTGCATACTCGGTAACCTGGTATCATTTTGCTGAA TTCAACTTATACTTGTTGATCAACAAAAACATGCAATTCATTTTGCTGAA	6450
TTCAACTTATACTTGTTGATCAACAAAAACATGCAATTCATTTTGGTGATAAAAAAAA	6500
	6550
GTCGGATGCTCTTTGCAACACATTACCACCCTCTCACCAAGGAATTCGCG	6600
TCTCACCCACGTGTCACCTCGAAACACATGGCTTGCGCATTCAAATCAAG	

Figure 11 (Continued)

ATCTGATTATCAACCACGTGGTTGTGATCAAGACCTAGTGTTCTTGTACC	6650
GTTTAACCGAGGGAGCTTGTCCTGAGAGCTACGGACTTCAAGTGGCACTC	6700
ATGGCTGGAATACCAAACCAAGTGGTTGAAACAGCATCAGGTGCTGCTCA	6750
AGCCATGAAGAGATCAATTGGGGAAAACTTCAAGTCAAG	6800
CTGAGTTCTCAAGTCTGCATGAAGACTGGCTCAAGTCATTGGTGGGTATT	6850
TCTCGAGTCGCCCACAACAATGCCCCCATTGGCGAAGATGACTACGACAC	6900
TTTGTTTTGCTTATGGCATGAGATCAAATCCTCTTACTGTGTTCCCAAAT	6950
AAATGGCTATGACATAACACTATCTGAAGCTCGTTAAGTCTTTTGCTTCT	7000
CTGATGTTTATTCCTCTTAAAAAATGCTTATATATCAAAAAATTGTTTCC	7050
TCGATTATAACAAGATTATATATGTATCTGTCGGTTTAGCTATGGTATAT	7100
AATATATGTATGTTCATGAGATTGGTCAAGAGAAATACTCACAAACAGTA	7150
TATTAAGAAGGAAATATGTTTATGCATTAATTTAAGTTTCAAGATAAACT	7200
GCAAATAACCTCGACTAAAGTTGCAAAGACCAAACACAAATTACAAAACT	7250
TATAAGACTTAAGTTCTGAATTCCCTAAAACCAAAAAAAA	7300
TATTTTGTTGCATCTACAAACACACAAACCTACATAGTTTATAACTTAC	7350
TCATCACTGAGATTAACATCAGAATCATTCTCCATTTCTTCATCTTCACT	7400
CTCATCATCACCACCACCATGATGATTCTCCTCCTCTTCACGTAACC	7450
TAGCAATCTCACTCTGAGCTCTATCAACAATCTGCTTCTTCTGCAACTCC	7500
AAATCTCTCTGAAAATCAGCTCTCATCTTCTCCAACTCCTTCATTTGCTC	7550
TTTCTTACTCTTCTCCATCTTCTCATAAACCTTCCCAAACCTCTCAACAG	7600
AATCCGCCAACATCTTATACGAAGCAGCGTCATTAACCTTCTTCCTCTCG	7650
TACTCAACCTCATCCTCATCCTCCTCCTCTCAGAATCACCAGGACT	7700
ATCCATCATCTCATCAAACCCATTAGACTTATCTAAATAAA	7750
TCATAAACACAAACTCACCTGAATCAACACCACAAGCTAAACCTAAATCC	7800
GACTTGGGCGAAACACAAAGCAACATATCCAACTTATTGAAAAACGACCA	7850
TTTACTTGAACCTAAACCTGATTTCTCAACCTTAATCTTCTCTTTTCTAT	7900
ACTTCCTCTTCAAGTCATCAATCATTCTCCTACATTGCGTCTCAGATTTC	7950
TCCATCCTTAGCTCCTCACTCTCTCAGCTACTTCATTCCAATCCTC	8000
GTTCCTCAAACTCCTTCTACCCAATTGCAAAAACCTATCTCCCCAAACTT	8050
CAAGCAACACAA	8062

Figure 11 (Continued)



Comments/References: AtMSH3 3' side antisense : AtMSH3 3' (13 = 2104bp) from pUC18/13 Sal1/Sst1/T4 into pCW164 BamH1/T4 in Agrobacterium LBA44O4




Figure 13

Comments/References: AtMSH6 (S8) 3' side antisens : 62 Sal1/Sst1/T4 (1379bp)

into pCW164 BamH1/T4




Figure 14

Comments/References: AtMSH6 3'/AtMSH3 3' antisense : AtMSH6 (S8) 3' side (62=1379bp) Sal1/Sst1/T4 into pPF13 (pCW164 AtMSH3 (S5) 3' side (13=2104) antisens)/Sma1. in LBA4404




Figure 15

Comments/References: AtMSH3 3'/AtMSH6 3' antisens (D): AtMSH3 (S5) 3' side (13=2104bp) Sal1/Sst1/T4 into pPF14 (AtMSH6 (S8) 3'side (62=1379bp) antisense into pCW164)/Sma1. in LBA4404




Figure 16

Comments/References: AtMSH3 (S8) complete, sense orientation : pPF26 (3342bp) Sma1 into pCW164 Sma1






Figure 17

Commente/References: pPZP111 with codA EcoR1 cassette in EcoR1 site and HInd3 GUS cassette in Hind3 site. KanR. All genes under Promoter/terminator 35S



## p3243



1

SEQUENCE LISTING

```
Rhone-Poulenc Agro; Betzner, Andreas Stefan; Doutriaux,
<110>
             Marie-Pascale; Freyssinet, Georges; Perez, Pascual.
            Methods for obtaining plant varieties
<120>
            395498C
<130>
<150>
             PO9745
<151>
             1997-10-10
<160>
             98
<210>
             1
<211>
             23
<212>
             DNA
<213>
             Artificial sequence
<220>
<221>
             modified_base
<222>
             11
<223>
             Ι
<220>
<221>
             modified_base
<222>
             14
<223>
             Ι
<220>
             modified_base
<221>
<222>
             17
<223>
<220>
             Degenerate oligonucleotides UPMU used to isolate AtMSH3 and
<223>
             AtMSH6.
<300>
<301>
             Reenan and Kolodner
<302>
             Genetics
<303>
             132
<306>
             963-973
<307>
             1992
<400>
             1
                                                                       23
ctggatccac nggnccnaay atg
```

<210> 2 <211> 23 <212> DNA

.. . . . .

2

```
Artificial sequence
<213>
<220>
<221>
             modified_base
<222>
             15
<223>
             Ι
<220>
<221>
             modified_base
<222>
             18
<223>
<220>
             Degenerate oligonucleotides DOMU used to isolate AtMSH3 and
<223>
             AtMSH6.
<300>
             Reenan and Kolodner
<301>
<302>
             Genetics
<303>
             132
             963-973
<306>
             1992
<307>
<400>
             2
                                                                       23
ctggatccrt artgngtnrc raa
<210>
             3
<211>
             24
<212>
             DNA
             Artificial sequence
<213>
<220>
             MSH3 specific primer 636 for PCR using cDNA of Arabidopsis
<223>
             thaliana ecotype Columbia
<400>
             3
                                                                       24
tgctagtgcc tcttgcaagc tcat
<210>
             4
             27
<211>
<212>
             DNA
<213>
             Artificial sequence
<220>
             Primer AP1 for PCR using cDNA of Arabidopsis thaliana ecotype
<223>
             Columbia containing adapter sequences ligated to both its
             ends
<400>
```

	3	
	amount at aggs	27
ccatcctaat a	cgactcact atagggc	
<210>	5	
<211>	23	
<212>	DNA	
<213>	Artificial sequence	
<220>		
<223>	Primer AP2 for PCR using cDNA of Arabidopsis thaliana ed	cotype
	Columbia containing adapter sequences ligated to both it	S
	ends	
<400>	5	
		23
actcactata g	ggctcgagc ggc	23
<210>	6	
<211>	30	
<212>	DNA	
	Artificial sequence	
<220>		
<223>	MSH3 specific primer S525 for PCR using cDNA of Arabido	psis
	thaliana ecotype Columbia	
<400>	6	
		30
aggttetgat t	atgtgtgac gctttactta	
<210>	7	
<211>	29	
<212>	DNA	
<213>	Artificial sequence	
<220>		
<223>	MSH3 specific primer S51 for PCR using cDNA of Arabidop	SIS
	thaliana ecotype Columbia	
<400>	7	
		29
ggatcgggta	ctgggttttg agtgtgagg	2,7
<210>	8	
<210>	24	
<212>	DNA	
<213>	Artificial sequence	
<220>		
<223>	MSH3 specific primer 635 for PCR using cDNA of Arabidor	sis
	thaliana ecotype Columbia	

4

< 400>	8
gcacgtgctt g	ratggtgttt tcac 24
55-5	
<210>	9
<211>	28
<212>	DNA
<213>	Artificial sequence
<220>	
<223>	MSH3 specific primer S523 for PCR using cDNA of Arabidopsis
(223)	thaliana ecotype Columbia
<400>	9
-	
tcagacagta t	ccagcatgg cagaagta 28
<210>	10
<211>	33
<212>	DNA
<213>	Artificial sequence
<220>	
<223>	MSH3 specific primer 1S5 for PCR using cDNA of Arabidopsis
	thaliana ecotype Columbia
<400>	10
atcccgggat g	gggcaagcaa aagcagcaga cga 33
<210>	11
<211>	27
<212>	DNA
	2
<213>	Artificial sequence
<220>	
<223>	MSH3 specific primer S53 for PCR using cDNA of Arabidopsis
	thaliana ecotype Columbia
<400>	11
gacaaagagc g	gaaatgaggc cccttgg 27
<210>	12
<211>	1250
<212>	DNA
<213>	Arabidopsis thaliana ecotype Columbia
<223>	Clone 52

< 400 >	12					
cccgggatgg	gcaagcaaaa	gcagcagacg	atttctcgtt	tcttcgctcc	caaacccaaa	60
teceegaete	acgaac <b>cg</b> aa	tccggtagcc	gaatcatcaa	caccgccacc	gaagatatcc	120
gccactgtat	ccttctctcc	ttccaagcgt	aagcttctct	ccgaccacct	cgccgccgcg	180
tcacccaaaa	agcctaaact	ttctcctcac	actcaaaacc	cagtacccga	tcccaattta	240
caccaaagat	ttctccagag	atttctggaa	ccctcgccgg	aggaatatgt	tcccgaaacg	300
tcatcatcga	ggaaatacac	accattggaa	cagcaagtgg	tggagctaaa	gagcaagtac	360
ccagatgtgg	ttttgatggt	ggaagttggt	tacaggtaca	gattcttcgg	agaagacgcg	420
gagatcgcag	cacgcgtgtt	gggtatttac	gctcatatgg	atcacaattt	catgacggcg	480
agtgtgccaa	catttcgatt	gaatttccat	gtgagaagac	tggtgaatgc	aggatacaag	540
attggtgtag	tgaagcagac	tgaaactgca	gccattaagt	cccatggtgc	aaaccggacc	600
ggcccttttt	tccggggact	gtcggcgttg	tataccaaag	ccacgcttga	agcggctgag	660
gatataagtg	gtggttgtgg	tggtgaagaa	ggttttggtt	cacagagtaa	tttcttggtt	720
tgtgttgtgg	atgagagagt	taagtcggag	acattaggct	gtggtattga	aatgagtttt	780
gatgttagag	tcggtgttgt	tggcgttgaa	atttcgacag	gtgaagttgt	ttatgaagag	840
ttcaatgata	atttcatgag	aagtggatta	gaggctgtga	ttttgagctt	gtcaccagct	900
gagctgttgc	ttggccagcc	tctttcacaa	caaactgaga	agtttttggt	ggcacatgct	960
ggacctacct	caaacgttcg	agtggaacgt	gcctcactgg	attgtttcag	caatggtaat	1020
gcagtagatg	aggttatttc	attatgtgaa	aaaatcagcg	caggtaactt	agaagatgat	1080
aaagaaatga	agctggaggc	tgctgaaaaa	ggaatgtctt	gcttgacagt	tcatacaatt	1140
atgaacatgc	cacatctgac	tgttcaagcc	ctcgccctaa	cgttttgcca	tctcaaacag	1200
tttggatttg	aaaggatcct	ttaccaaggg	gcctcatttc	gctctttgtc		1250

<210> 13 <211> 34 <212> DNA

<213> Artificial sequence

<220>

MSH3 specific primer 2S5 for PCR using cDNA of Arabidopsis <223> thaliana ecotype Columbia

PCT/EP98/06977

			6			•
< 400 >	13					
atcccgggtc	aaaatgaaca	agttggtttt	agtc			34
<210><211>	14 27					
<212> <213>	DNA Artificia	l seguence				
(213)	ALCITICIA	1 Dequence				
<220>						
<223>	_	ific primer ecotype Col		R using cDN	IA of Arabido	psis
<400>	14					
gccacatctg	actgttcaag	ccctcgc				27
<210>	15					
<211>	2110					
<212>	DNA					
<213>	Arabidops	is thaliana	ecotype Co	olumbia		
<223>	Clone 13					
<400>	15					
gccacatctg	actgttcaag	ccctcgccct	aacgttttgc	catctcaaac	agtttggatt	60
tgaaaggatc	ctttaccaag	gggcctcatt	tcgctctttg	tcaagtaaca	cagagatgac	120
tctctcagcc	aatactctgc	aacagttgga	ggttgtgaaa	aataattcag	atggatcgga	180
atctggctcc	ttattccata	atatgaatca	cacacttaca	gtatatggtt	ccaggcttct	240
tagacactgg	gtgactcatc	ctctatgcga	tagaaatttg	atatctgctc	ggcttgatgc	300
tgtttctgag	atttctgctt	gcatgggatc	tcatagttct	tcccagctca	gcagtgagtt	360
ggttgaagaa	ggttctgaga	gagcaattgt	atcacctgag	ttttatctcg	tgctctcctc	420
agtcttgaca	gctatgtcta	gatcatctga	tattcaacgt	ggaataacaa	gaatctttca	480
tcggactgct	aaagccacag	agttcattgc	agttatggaa	gctattttac	ttgcggggaa	540
gcaaattcag	cggcttggca	taaagcaaga	ctctgaaatg	aggagtatgc	aatctgcaac	600
tgtgcgatct	actcttttga	gaaaattgat	ttctgttatt	tcatcccctg	ttgtggttga	660
caatgccgga	aaacttctct	ctgccctaaa	taaggaagcg	gctgttcgag	gtgacttgct	720
cgacatacta	atcacttcca	gcgaccaatt	tcctgagctt	gctgaagctc	gccaagcagt	780

trtagtcatc agggaaaagc tggattcctc gatagcttca tttcgcaaga agctcgctat 840

togaaatttg gaatttotto	aagtgtcggg	gatcacacat	ttgatagagc	tgcccgttga	900
ttccaaggtc cctatgaatt	gggtgaaagt	aaatagcacc	aagaagacta	ttcgatatca	960
tcccccagaa atagtagctg	gcttggatga	gctagctcta	gcaactgaac	atcttgccat	1020
tgtgaaccga gcttcgtggg	atagtttcct	caagagtttc	agtagatact	acacagattt	1080
taaggctgcc gttcaagctc	ttgctgcact	ggactgtttg	cactcccttt	caactctatc	1140
tagaaacaag aactatgtcc	gtcccgagtt	tgtggatgac	tgtgaaccag	ttgagataaa	1200
catacagtct ggtcgtcatc	ctgtactgga	gactatatta	caagataact	tcgtcccaaa	1260
tgacacaatt ttgcatgcag	aaggggaata	ttgccaaatt	atcaccggac	ctaacatggg	1320
aggaaagagc tgctatatcc	gtcaagttgc	tttaatttcc	ataatggctc	aggttggttc	1380
ctttgtacca gcgtcattcg	ccaagctgca	cgtgcttgat	ggtgttttca	ctcggatggg	1440
tgcttcagac agtatccagc	atggcagaag	tacctttcta	gaagaattaa	gtgaagcgtc	1500
acacataatc agaacctgtt	cttctcgttc	gcttgttata	ttagatgagc	ttggaagagg	1560
cactagcaca cacgacggtg	tagccattgc	ctatgcaaca	ttacagcatc	tcctagcaga	1620
aaagagatgt ttggttcttt	ttgtcacgca	ttaccctgaa	atagctgaga	tcagtaacgg	1680
attcccaggt tctgttggga	cataccatgt	ctcgtatctg	acattgcaga	aggataaagg	1740
cagttatgat catgatgatg	tgacctacct	atataagctt	gtgcgtggtc	tttgcagcag	1800
gagetttggt tttaaggttg	ctcagcttgc	ccagatacct	ccatcatgta	tacgtcgagc	1860
catttcaatg gctgcaaaat	tggaagctga	ggtacgtgca	agagagagaa	atacacgcat	1920
gggagaacca gaaggacatg	aagaaccgag	aggcgcagaa	gaatctattt	cggctctagg	1980
tgacttgttt gcagacctga	aatttgctct	ctctgaagag	gacccttgga	aagcattcga	2040
gtttttaaag catgcttgga	agattgctgg	caaaatcaga	ctaaaaccaa	cttgttcatt	2100
ttgacccggg		•			2110

<210> 16

<211> 29

<212> DNA

<213> Artificial sequence

<220>

<223> MSH3 specific primer S51 for PCR using cDNA of Arabidopsis
thaliana ecotype Columbia

<400> 16

ggat	.cggg	rta c	tggg	rttt	g ag	tgtg	agg								•	29
<210	<b> </b> >		17													
<211			30													
<212				DNA												
<213					ial	SAGU	ence	,								
\21J	,		MI (	.1110	.141	sequ	ence	•								
<220	>															
<223	>		MSH	3 sp	ecif	ic p	rime	r S5	25 £	or P	CR u	sing	CDN	IA of	Arab	idopsis
			tha	lian	a ec	otyp	e Co	lumb	ia							
<400	>		17													
aggt	tctg	rat t	atgt	gtga	ac go	ttta	ictta	ı								30
<210	) >		18													
<211	.> '		352	2												
<212	: >		DNA													
<213	<b>&gt;</b>		Ara	bido	psis	tha	lian	a ec	otyp	e Co	lumi	oia				
226																
<220			CDC	,												
<221			CDS		, ,											
<222			•		(3							•			- 1-	
<223	S >						ıgtn	CDNA	anc	ı aec	iuceo	sec	luenc	e or	the	encoded
			po1	yper	otide	•										
<400	) >		18													
ccta	agaa	ag o	gege	gaaa	aa tt	ggca	aacco	aag	gttcg	gcca	tago	cac	gac o	cacga	accttc	60
catt	tct	ett a	aaac	ggagg	ga ga	attac	gaat	: aaa	agcaa	act						99
atq	ggc	aaq	caa	aaq	caq	caq	acq	att	tct	cat	ttc	ttc	qct	ccc	aaa	147
										_				Pro		
1		•		5					10	_		- " -		15	•	
ccc	aaa	tcc	cca	act	cac	gaa	cca	aat	cca	gta	acc	gaa	tca	tca	aca	195
									_	_	_	_		Ser		
	•		20				_	25					30		-	
cca	cca	cca	aaq	ata	tcc	acc	act	qta	tcc	ttc	tct	cct	tcc	aag	cat	243
			_			_		_						Lys	_	
		35	-1-				40					45		_,_	•••-	
ааσ	CEE	ctc	ticc	gac	cac	ctc	acc	acc	aca	tca	CCC	222	aad	cct	222	291
														Pro		2,7
-10	50	u			****	55	774	ALG	ALG	JEI	60	Lys	دوب	110	د ر د	
	20					J.)					90					
ctt	tct	CCF	cac	act	caa	220	CC3	at >	~~~	are t	ccc	aat	tta	cac	caa	339
								_		_				His		332
שפע		0			70			val	FLO	ASP	LLU	W211	<b></b> u		0.0	

									-							
								ccc Pro								387
								aca Thr 105								435
								gtg Val								483
								gac Asp								531
								cac His								579
								gtg Val								627
								act Thr 185								675
								ttt Phe								723
								gct Ala								771
ggt Gly 225	gg:t Gly	gaa Glu	gaa Glu	ggt Gly	ttt Phe 230	ggt Gly	tca Ser	cag Gln	agt Ser	aat Asn 235	ttc Phe	ttg Leu	gtt Val	tgt Cys	gtt Val 240	819
gtg Val	gat Asp	gag Glu	aga Arg	gtt Val 245	aag Lys	tcg Ser	gag Glu	aca Thr	tta Leu 250	ggc Gly	tgt Cys	ggt Gly	att Ile	gaa Glu 255	atg Met	867
								gtt Val 265								915
gaa Glu	gtt Val	gtt Val 275	tat Tyr	gaa Glu	gag Glu	ttc Phe	aat Asn 280	gat Asp	aat Asn	ttc Phe	atg Met	aga Arg 285	agt Ser	gga Gly	tta Leu	963

													ggc Gly		1011
					_	_		_		_		-	gga Gly		1059
		-	_			_	_		_	_	_		agc Ser 335		1107
_		_	_										agc Ser		1155
		_	_	_					-		_		gaa Glu		1203
	_	-	-						-				cat His		1251
	_	_		-		_		_				_	ttt Phe		1299
	_						-			-		_	tca Ser 415	_	1347
		 _				Ala			_		_	_	gag Glu	-	1395
		Asn											cat His		1443
													cac His		1491
	Thr												ctt Leu		1539
							_				_		tcc Ser 495	-	1587

													att Ile 510			16	335
												-	atg Met			16	83
													cgg <b>Ar</b> g			17	731
													ctt Leu			17	779
													atg Met			18	27
													ttg Leu 590			18	75
													ctt Leu			19	23
													gac Asp			19	71
													cgc Arg			20	19
													tca Ser			20	67
aag Lys	aag Lys	ctc Leu	gct Ala 660	att Ile	cga Arg	aat Asn	ttg Leu	gaa Glu 665	ttt Phe	ctt Leu	caa Gln	gtg Val	tcg Ser 670	gly aaa	atc Ile	21	15
aca Thr	cat His	ttg Leu 675	ata Ile	gag Glu	ctg Leu	ccc Pro	gtt Val 680	gat Asp	tcc Ser	aag Lys	gtc Val	cct Pro 685	atg His	aat Asn	tgg Trp	21	63
													ccc Pro			22	11

												ctt Leu		2259
												agt Ser 735		2307
					_	-		_		-	_	ctg Leu	_	2355
_	_						_		_			gtc Val	-	2403
			_	_	_		_					cag Gln		2451
												gtc Val		2499
												atc Ile 815		2547
						-			_		-	gct Ala		2595
												ttc Phe		2643
												tca Ser		2691
												gaa Glu		2739
												tta Leu 895		2787
												gcc Ala		2835

														ctt Leu		2883
_	_				-		-			_				cca Pro		2931
	_								-		_			gat Asp		2979
	_		_											gtg Val 975		3027
			_						_	-	-			gcc Ala		3075
						Arg					Met			aaa Lys		3123
Glu					Ala					Thr				gaa Glu		3171
	Gly			Glu					Glu					gct Ala		3219
			Phe					Phe		Leu			Glu	gac Asp 1055		3267
		Ala		-			Lys		-		_	Ile		ggc		3315
	Arg				act Thr	Cys			tga	ttta	atc '	ttaa	catt	at		3362
agca	aact	gca :	aggt	cttg	at ca	atct	gtta	g tt	gcgt	acta	act	tatg	tgt a	atta	gtataa	3422
caa	gaaa	aga (	gaat	taga	ga ga	atgg	attc	t aa	tccg	gtgt	tgc	agta	cat (	cttt	tctcca	3482
ccc	gcat	aaa	aaaa	aaaa	aa aa	aaaa	aaaa	a aa	aaaa	aaaa						3522

<210> 19 <211> 1081 <212> PRT

<213 <223					_	tha MSH		a ec	otyp	e Co	lumb	ia			
< 400	) >		19												
Met 1	Gly	Lys	Gln	Lys 5	Gln	Gln	Thr	Ile	Ser 10	Arg	Phe	Phe	Ala	Pro 15	Lys
Pro	Lys	Ser	Pro 20	Thr	His	Glu	Pro	Asn 25	Pro	Val	Ala	Glu	Ser 30	Ser	Thr
Pro	Pro	Pro 35	Lys	Ile	Ser	Ala	Thr 40	Val	Ser	Phe	Ser	Pro 45	Ser	Lys	Arg
Lys	Leu 50	Leu	Ser	Asp	His	Leu 55	Ala	Ala	Ala	Ser	Pro 60	Lys	Lys	Pro	Lys
Leu 65	Ser	Pro	His	Thr	Gln 70	Asn	Pro	Val	Pro	Asp 75	Pro	Asn	Leu	His	Gln 80
Arg	Phe	Leu	Gln	Arg 85	Phe	Leu	Glu	Pro	Ser 90	Pro	Glu	Glu	Tyr	Val 95	Pro
Glu	Thr	Ser	Ser 100	Ser	Arg	Lys	Tyr	Thr 105	Pro	Leu	Glu	Gln	Gln 110	Val	Val
Glu	Leu	Lys 115	Ser	Lys	Tyr	Pro	Asp 120	Val	Val	Leu	Met	Val 125	Glu	Val	Gly
Tyr	Arg 130	Tyr	Arg	Phe	Phe	Gly 135	Glu	Asp	Ala	Glu	Ile 140	Ala	Ala	Arg	Val
Leu 145	Gly	Ile	Tyr	Ala	His 150	Met	Asp	His	Asn	Phe 155	Met	Thr	Ala	Ser	Val 160
Pro	Thr	Phe	_	Leu 165	Asn	Phe	His		Arg 170	_	Leu	Val	Asn	Ala 175	Gly
Tyr	Lys	Ile	Gly 180	Val	Val	Lys	Gln	Thr 185	Glu	Thr	Ala	Ala	Ile 190	Lys	Ser
His	Gly	Ala 195	Asn	Arg	Thr	Gly	Pro 200	Phe	Phe	Arg	Gly	Leu 205	Ser	Ala	Leu
Tyr	Thr 210	Lys	Ala	Thr	Leu	Glu 215	Ala	Ala	Glu	Asp	Ile 220	Ser	Gly	Gly	Cys
Gly 225	Gly	Glu	Glu	Gly	Phe 230	Gly	Ser	Gln	Ser	Asn 235	Phe	Leu	Val	Cys	Val 240
Val	Asp	Glu	Arg	Val 245	Lys	Ser	Glu	Thr	Leu 250	Gly	Cys	Gly	Ile	Glu 255	Met

Ser Phe Asp Val Arg Val Gly Val Gly Val Glu Ile Ser Thr Gly Glu Val Val Tyr Glu Glu Phe Asn Asp Asn Phe Met Arg Ser Gly Leu Glu Ala Val Ile Leu Ser Leu Ser Pro Ala Glu Leu Leu Gly Gln Pro Leu Ser Gln Gln Thr Glu Lys Phe Leu Val Ala Met Ala Gly Pro Thr Ser Asn Val Arg Val Glu Arg Ala Ser Leu Asp Cys Phe Ser Asn Gly Asn Ala Val Asp Glu Val Ile Ser Leu Cys Glu Lys Ile Ser Ala Gly Asn Leu Glu Asp Asp Lys Glu Met Lys Leu Glu Ala Ala Glu Lys Gly Met Ser Cys Leu Thr Val His Thr Ile Met Asn Met Pro His Leu Thr Val Gln Ala Leu Ala Leu Thr Phe Cys His Leu Lys Gln Phe Gly Phe Glu Arg Ile Leu Tyr Gln Gly Ala Ser Phe Arg Ser Leu Ser Ser Asn Thr Glu Met Thr Leu Ser Ala Asn Thr Leu Gln Gln Leu Glu Val Val Lys Asn Asn Ser Asp Gly Ser Glu Ser Gly Ser Leu Phe His Asn Met Asn His Thr Leu Thr Val Tyr Gly Ser Arg Leu Leu Arg His Trp Val Thr His Pro Leu Cys Asp Arg Asn Leu Ile Ser Ala Arg Leu Asp Ala Val Ser Glu Ile Ser Ala Cys Met Gly Ser His Ser Ser Ser Gln Leu Ser Ser Glu Leu Val Glu Glu Gly Ser Glu Arg Ala Ile Val Ser Pro Glu Phe Tyr Leu Val Leu Ser Ser Val Leu Thr Ala Met Ser Arg Ser Ser Asp Ile Gln Arg Gly Ile Thr Arg Ile Phe His Arg Thr Ala

Lys 545	Ala	Thr	Glu	Phe	11e 550	Ala	Val	Met	Glu	Ala 555	Ile	Leu	Leu	Ala	Gl <sub>3</sub> 560
Lys	Gln	Ile	Gln	Arg 565	Leu	Gly	Ile	Lys	Gln 570	Asp	Ser	Glu	Met	Arg 575	Ser
Met	Gln	Ser	Ala 580	Thr	Val	Arg	Ser	Thr 585	Leu	Leu	Arg	Lys	Leu 590	Ile	Sei
Val	Ile	Ser 595	Ser	Pro	Val	Val	Val 600	Asp	Asn	Ala	Gly	Lys 605	Leu	Leu	Ser
Ala	Leu 610	Asn	Lys	Glu	Ala	Ala 615	Val	Arg	Gly	Asp	Leu 620	Leu	Asp	Ile	Leu
Ile 625	Thr	Ser	Ser	Asp	Gln 630	Phe	Pro	Glu	Leu	Ala 635	Glu	Ala	Arg	Gln	Ala 640
Val	Leu	Val	Ile	Arg 645	Glu	Lys	Leu	Asp	Ser 650	Ser	Ile	Ala	Ser	Phe 655	Arg
Lys	Lys	Leu	Ala 660	Ile	Arg	Asn	Leu	Glu 665	Phe	Leu	Gln	Val	Ser 670	Gly	Ile
Thr	His	Leu 675	Ile	Glu	Leu	Pro	Val 680	Asp	Ser	Lys	Val	Pro 685	His	Asn	Trp
Val	Lys 690	Val	Asn	Ser	Thr	Lys 695	Lys	Thr	Ile	Arg	Tyr 700	His	Pro	Pro	Glu
Ile 705	Val	Ala	Gly	Leu	Asp 710	Glu	Leu	Ala	Leu	Ala 715	Thr	Glu	His	Leu	Ala 720
Ile	Val	Asn	Arg	Ala 725	Ser	Trp	Asp	Ser	Phe 730	Leu	Lys	Ser	Phe	Ser 735	Arg
Tyr	Tyr	Thr	Asp 740	Phe	Lys	Ala	Ala	Val 745	Gln	Ala	Leu	Ala	Ala 750	Leu	Asp
Cys	Leu	His 755	Ser	Leu	Ser	Thr	Leu 760	Ser	Arg	Asn	Lys	Asn 765	Tyr	Val	Arg
Pro	Glu 770	Phe	Val	Asp	Asp	Cys 775	Glu	Pro	Val	Glu	Ile 780	Asn	Ile	Gln	Ser
Gly 785	Arg	His	Pro	Val	Leu 790	Glu	Thr	Ile	Leu	Gln 795	Asp	Asn	Phe	Val	Pro 800
Asn	Asp	Thr	Ile	Leu 805	His	Ala	Glu	Gly	Glu 810	Tyr	Cys	Gln	Ile	Ile 815	Thr
Gly	Pro	Asn	Met 820	Gly	Gly	Lys	Ser	Cys	Tyr	Ile	Arg	Gln	Val	Ala	Lev

Ile		Ile 835	Met	Ala	Gln		Gly 840	Ser	Phe	Val	Pro	Ala 845	Ser	Phe	Ala
_	Leu 850	His	Val	Leu		Gly 855	Val	Phe	Thr		Met 860	Gly	Ala	Ser	Asp
Ser 865	Ile	Gln	His	Gly	Arg 870	Ser	Thr	Phe	Leu	Glu 875	Glu	Leu	Ser	Glu	Ala 880
Ser	His	Ile	Ile	Arg 885	Thr	Cys	Ser	Ser	Arg 890	Ser	Leu	Val	Ile	Leu 895	Asp
Glu	Leu	Gly	Arg	Gly	Thr	Ser	Thr	His 905	Asp	Gly	Val	Ala	Ile 910	Ala	Tyr
Ala	Thr	Leu 915	Gln	His	Leu	Leu	Ala 920	Glu	Lys	Arg	Cys	Leu 925	Val	Leu	Phe
Val	Thr 930	His	Tyr	Pro	Glu	Ile 935	Ala	Glu	Ile	Ser	Asn 940	Gly	Phe	Pro	Gly
Ser 945	Val	Gly	Thr	Tyr	His 950	Val	Ser	Tyr	Leu	Thr 955	Leu	Gln	Lys	qsA	Lys 960
Gly	Ser	Tyr	Asp	His 965	Asp	Asp	Val	Thr	Tyr 970	Leu	Tyr	Lys	Leu	Val 975	Arg
Gly	Leu	Cys	Ser 980	Arg	Ser	Phe	Gly	Phe 985	Lys	Val	Ala	Gln	Leu 990	Ala	Gln
lle	Pro	Pro 995	Ser	Cys	Île		Arg 1000	Ala	Ile	Ser		Ala 1005	Ala	Lys	Leu
	Ala 1010	Glu	Val	Arg		Arg 1015		Arg	Asn		Arg 1020		Gly	Glu	Pro
Glu 102	-	His	Glu		Pro 1030	Arg	Gly	Ala	Glu	. Glu 1035		Ile	Ser		Leu 1040
Gly	Asp	Leu		Ala 1045		Leu	Lys		Ala 1050		Ser	Glu		Asp 1055	Pro
Trp	Lys	Ala	Phe		Phe	Leu	Lys	His 1065		Trp	Lys	Ile	Ala 1070		Lys
Ile	Arg	Leu 1075	-	Pro	Thr	Cys	Ser 1080		:						
<21	0 >		20	ı											

<210> 20 <211> 24 <212> DNA <213> Artificial sequence WO 99/19492 PCT/EP98/06977

18

<220>		
<223>	MSH6 specific primer 638 for PCR using cDNA of Arabidopsis thaliana ecotype Columbia	
<400>	20	
tctctaccag c	gtgacgaaaa accg 2	4
<210>	21	
<211>	28	
<212>	DNA	
<213>	Artificial sequence	
<220>		
<223>	Primer S81 for PCR using cDNA of Arabidopsis thaliana ecoty Columbia	pe
<400>	21	
cgtcgccttt a	agcatecect teetteae 2	8.
<210>	22	
<211>	30	
<212>	DNA	
<213>	Artificial sequence	
<220>		
<223>	MSH6 specific primer S823 for PCR using cDNA of Arabidopsis thaliana ecotype Columbia	
<400>	22	
gcttggcgca (	cctaatagaa tcatgacagg 3	0
<210>	23	
<211>	24	
<212>	DNA	
<213>	Artificial sequence	
<220>		
<223>	MSH6 specific primer 637 for PCR using cDNA of Arabidopsis	
	thaliana ecotype Columbia	
<400>	23	
gacagcgtca	gttcttcaga atgc 2	24
<210>	24	
<211>	33	

<212> DNA

<213>	Artificial sequence	
<220>		
<223>	MCUC anocific number 100 for DCD union apply of Archidonai	_
.2237	MSH6 specific primer 1S8 for PCR using cDNA of Arabidopsi thaliana ecotype Columbia	5
	charrana ecocype corumbia	
<400>	24	
atcccgggat	gcagcgccag agatcgattt tgt	33
<210>	25	
<211>	27	
<212>	DNA	
<213>	Artificial sequence	
<220>		
<223>	MSH6 specific primer S83 for PCR using cDNA of Arabidopsi	5
	thaliana ecotype Columbia	
<400>	25	
cgctatctat	ggctgcttcg aattgag	27
.010		
	26	
	1385	
	DNA	
	Arabidopsis thaliana ecotype Columbia	
<223>	Clone 43	
<400>	26	
	_	
cccgggatgc	agegecagag ategatiting tettiettee aaaaaeeeae ggeggegaet	60
accaaccatt	tagtitagga ogstgotgot saggaggaga gaggasgaga saggas	120
acgaagggcc	tggtttccgg cgatgctgct agcggcgggg gcggcagcgg aggaccacga	120
tttaatgtga	aggaagggga tgctaaaggc gacgcttctg tacgttttgc tgtttcgaaa	180
,		
tctgtcgatg	aggttagagg aacggatact ccaccggaga aggttccgcg tcgtgtcctg	240
ccgtctggat	ttaageegge tgaateegee ggtgatgett egteeetgtt etceaatatt	300
atgcataagt	ttgtaaaagt cgatgatcga gattgttctg gagagaggag ccgagaagat	360
		300
gttgttccgc	tgaatgattc atctctatgt atgaaggcta atgatgttat tcctcaattt	420
cgttccaata	atggtaaaac tcaagaaaga aaccatgctt ttagtttcag tgggagagct	480
gaacttagat	cagtagaaga tataggagta gatggcgatg ttcctggtcc agaaacacca	540
gggatgcgtc	cacgtgette tegettgaag egagttetgg aggatgaaat gaettttaag	600
	accordance of the second secon	300
gaggataagg	ticcidiati quacticiaec assangeida assingtica gostooget	<i>c</i>

tgtgg	agaga	agaaagaagt	aaacgaagga	accaaatttg	aatggcttga	gtcttctcga	720
atcag	ggatg	ccaatagaag	acgtcctgat	gatccccttt	acgatagaaa	gaccttacac	780
atacc	acctg	atgttttcaa	gaaaatgtct	gcatcacaaa	agcaatattg	gagtgttaag	840
agtga	atata	tggacattgt	gcttttcttt	aaagtgggga	aattttatga	gctgtatgag	900
ctaga	tgcgg	aattaggtca	caaggagctt	gactggaaga	tgaccatgag	tggtgtggga	960
aaatg	cagac	aggttggtat	ctctgaaagt	gggatagatg	aggcagtgca	aaagctatta	1020
gctcg	tggat	ataaagttgg	acgaatcgag	cagctagaaa	catctgacca	agcaaaagcc	1080
agagg	tgcta	atactataat	tccaaggaag	ctagttcagg	tattaactcc	atcaacagca	1140
agcga	gggaa	acatcgggcc	tgatgccgtc	catcttcttg	ctataaaaga	gatcaaaatg	1200
gagct	acaaa	agtgttcaac	tgtgtatgga	tttgcttttg	ttgactgtgc	tgccttgagg	1250
ttttg	ggttg	ggtccatcag	cgatgatgca	tcatgtgctg	ctcttggagc	gttattgatg	1320
caggt	ttctc	caaaggaagt	gttatatgac	agtaaagggc	tatcaagaga	agcacaaaag	1380
gctct	aagga	aatatacgtt	gacagggtct	acggcggtac	agttggctcc	agtaccacaa	1440
gtaat	ggggg	atacagatgc	tgctggagtt	agaaatataa	tagaatctaa	cggatacttt	1500
aaagg	ttott	ctgaatcatg	gaactgtgct	gttgatggtc	taaatgaatg	tgatgttgcc	1560
cttag	tgctc	ttggagagct	aattaatcat	ctgtctaggc	taaagctaga	agatgtactt	1620
aagca	rgggg	atattttcc	ataccaagtt	tacaggggtt	gtctcagaat	tgatggccag	1680
acgat	ggtaa	atcttgagat	atttaacaat	agctgtgatg	gtggtccttc	agggaccttg	1740
tacaa	atatc	ttgataactg	tgttagtcca	actggtaagc	gactcttaag	gaattggatc	1800
tgcca	tccac	tcaaagatgt	agaaagcatc	aataaacggc	ttgatgtagt	tgaagaattc	1860
acggc	aaact	cagaaagtat	gcaaatcact	ggccagtatc	tccacaaact	tccagactta	1920
gaaag	actgc	teggaegeat	caagtctagc	gttcgatcat	cagcctctgt	gttgcctgct	1980
cttct	gggga	aaaaagtgct	gaaacaacga	gttaaagcat	ttgggcaaat	tgtgaaaggg	2040
ttcag	aagtg	gaattgatct	gttgttggct	ctacagaagg	aatcaaatat	gatgagtttg	2100
cttta	taaac	tctgtaaact	tcctatatta	gtaggaaaaa	gcgggctaga	gttatttctt	2160
tctca	attcg	aagcagccat	agatagcg				2188

<210>

<211><211><212><213><223>	1385 DNA Arabidops Clone 62	sis thaliana	a ecotype Co	olumbia		
<400>	27					
catcagcctc	tgtgttgcct	gctcttctgg	ggaaaaaagt	gctgaaacaa	cgagttaaag	60
catttgggca	aattgtgaaa	gggttcagaa	gtggaattga	tctgttgttg	gctctacaga	120
aggaatcaaa	tatgatgagt	ttgctttata	aactctgtaa	acttcctata	ttagtaggaa	180
aaagcgggct	agagttattt	ctttctcaat	tcgaagcagc	catagatagc	gactttccaa	240
attatcagaa	ccaagatgtg	acagatgaaa	acgctgaaac	tctcacaata	cttatcgaac	300
tttttatcga	aagagcaact	caatggtctg	aggtcattca	caccataagc	tgcctagatg	360
tcctgagatc	ttttgcaatc	gcagcaagtc	tetetgetgg	aagcatggcc	aggcctgtta	420
tttttcccga	atcagaagct	acagatcaga	atcagaaaac	aaaagggcca	atacttaaaa	480
tccaaggact	atggcatcca	tttgcagttg	cagccgatgg	tcaattgcct	gttccgaatg	540
atatactcct	tggcgaggct	agaagaagca	gtggcagcat	tcatcctcgg	tcattgttac	600
tgacgggacc	aaacatgggc	ggaaaatcaa	ctcttcttcg	tgcaacatgt	ctggccgtta	660
tctttgccca	acttggctgc	tacgtgccgt	gtgagtcttg	cgaaatctcc	ctcgtggata	720
ctatcttcac	aaggcttggc	gcatctgata	gaatcatgac	aggagagagt	acctttttgg	780
tagaatgcac	tgagacagcg	tcagttcttc	agaatgcaac	tcaggattca	ctagtaatcc	840
ttgacgaact	gggcagagga	actagtactt	tcgatggata	cgccattgca	tactcggttt	900
ttcgtcacct	ggtagagaaa	gttcaatgtc	ggatgctctt	tgcaacacat	taccaccctc	960
tcaccaagga	attegegtet	cacccacgtg	tcacctcgaa	acacatggct	tgcgcattca	1020
aatcaagatc	tgattatcaa	ccacgtggtt	gtgatcaaga	cctagtgttc	ttgtaccgtt	1080
taaccgaggg	agcttgtcct	gagagctacg	gacttcaagt	ggcactcatg	gctggaatac	1140
caaaccaagt	ggttgaaaca	gcatcaggtg	ctgctcaagc	catgaagaga	tcaattgggg	1200
aaaacttcaa	gtcaagtgag	ctaagatctg	agttctcaag	tctgcatgaa	gactggctca	1260
agtcattggt	gggtatttct	cgagtcgccc	acaacaatgc	cccattggc	gaagatgact	1320
acgacacttt	gttttgctta	tggcatgaga	tcaaatcctc	ttactgtgtt	cccaaataac	1380

ccggg	1385
<210>	28
<211>	34
<212>	DNA
<213>	Artificial sequence
<220>	
<223>	MSH6 specific primer 2S8 for PCR using cDNA of Arabidopsis thaliana ecotype Columbia
<400>	26
atcccgggtt	atttgggaac acagtaagag gatt 34
<210>	29
<211>	27
<212>	DNA
<213>	Artificial sequence
<220>	
<223>	MSH6 specific primer S82 for PCR using cDNA of Arabidopsis thaliana ecotype Columbia
<400>	29
gcgttcgatc	atcagcctct gtgttgc 27
<210>	30
<211>	3606
<212>	DNA
<213>	Arabidopsis thaliana ecotype Columbia
<220>	
<221>	CDS
<222>	(142) (3468)
<223>	AtMSH6 full-length cDNA and deduced sequence of the encoded polypeptide
<400>	30
aaaagttgag	ccctgaggag tatcgtttcc gccatttcta cgacgcaagg cgaaaatttt 60
tggcgccaat	ctttccccc tttcgaattc tctcagctca aaacatcgtt tctctctcac 120
tctctctcac	aattccaaaa a atg cag cgc cag aga tcg att ttg tct ttc 171  Met Gln Arg Gln Arg Ser Ile Leu Ser Phe  1 5 10

							25							
												ggc Gly 25		219
									-			gtg Val		267
												tcg Ser	٠	315
_	_		_	-								gtt Val		363
												gst Gly		411
												gtc Val 105		459
												ccg Pro		507
_												caa Gln		555
			_									agt Ser		603
	_	_	-			-	_	_			_	gat Asp		651
								-		_	-	tct Ser 185		699
_	-	-	_	-		-			_		-	aag Lys		747
												ccg Pro		795

tgt Cys	gga Gly 220	gag Glu	aag Lys	aaa Lys	gaa Glu	gta Val 225	aac Asn	gaa Glu	gga Gly	acc Thr	aaa Lys 230	ttt Phe	gaa Glu	tgg Trp	ctt Leu	843
gag Glu 235	tct Ser	tc: Ser	cga Arg	atc Ile	agg Arg 240	gat Asp	gcc Ala	aat Asn	aga Arg	aga Arg 245	cgt Arg	cct Pro	gat Asp	gat Asp	ccc Pro 250	891
ctt Leu	tac Tyr	gat Asp	aga Arg	aag Lys 255	acc Thr	tta Leu	cac His	ata Ile	cca Pro 260	cct Pro	gat Asp	gtt Val	ttc Phe	aag Lys 265	aaa Lys	939
atg Met	tct Ser	gca Ala	tca Ser 270	caa Gln	aag Lys	caa Gln	tat Tyr	tgg Trp 275	agt Ser	gtt Val	aag Lys	agt Ser	gaa Glu 280	tat Tyr	atg Met	987
			ctt Leu													1035
			gaa Glu													1083
agt Ser 315	ggt Gly	gtg Val	gga Gly	aaa Lys	tgc Cys 320	aga Arg	cag Gln	gtt Val	ggt Gly	atc Ile 325	tct Ser	gaa Glu	agt Ser	GJÀ aaa	ata Ile 330	1131
			gtg Val													1179
			cta Leu 350						Ala						aat Asn	1227
act Thr	ata Ile	att Ile 365		agg Arg	aag Lys	cta Leu	gtt Val 370	Gln	gta Val	tta Leu	act Thr	cca Pro 375	Ser	aca Thr	gca Ala	1275
		Gly					Asp					Leu			aaa Lys	1323
	Ile					Gln					Val				gct Ala 410	1371
					Ala					Val					gat Asp	1419

								•	45							. ,
gat Asp	gca Ala	tca Ser	tgt Cys 430	gct Ala	gct Ala	ctt Leu	gga Gly	gcg Ala 435	tta Leu	ttg Leu	atg Met	cag Gln	gtt Val 440	tct Ser	cca Pro	1467
aag Lys	gaa Glu	gtg Val 445	tta Leu	tat Tyr	gac Asp	agt Ser	aaa Lys 450	ggg Gly	cta Leu	tca Ser	aga Arg	gaa Glu 455	gca Ala	caa Gln	aag Lys	1515
gct Ala	cta Leu 460	agg Arg	aaa Lys	tat Tyr	acg Thr	ttg Leu 465	aca Thr	ggg Gly	tct Ser	acg Thr	gcg Ala 470	gta Val	cag Gln	ttg Leu	gct Ala	1563
					atg Met 480											1611
					gga Gly											1659
tgt Cys	gct Ala	gtt Val	gat Asp 510	ggt Gly	cta Leu	aat Asn	gaa Glu	tgt Cys 515	gat Asp	gtt Val	gcc Ala	ctt Leu	agt Ser 520	gct Ala	ctt Leu	1707
					cat His											1755
					ttt Phe							Gly				1803
att Ile 555	gat Asp	ggc Gly	cag Gln	acg Thr	atg Met 560	gta Val	aat Asn	ctt Leu	gag Glu	ata Ile 565	Phe	aac Asn	aat Asn	agc Ser	tgt Cys 570	1851
					gly ggg					Tyr					Val	1899
				Lys	cga Arg				Asn					Pro		1947
			Glu		atc : Ile			Arg					Glu			1995
		Asn			agt Ser		Glr					ı Tyr				2043

			_	aga Arg 640	-					_		-			2091
				ttg Leu		_						-			2139
_	_		-	ttt Phe	_										2187
_	_	_	_	gct Ala		_	_	_			_	_	_	_	2235
			_	aaa Lys					-			_			2283
				caa Gln 720		_	_	_		_	_	_			2331
	_			gat Asp			-	-		_	_				2379
				ttt Phe											2427
			_	tgc Cys		-	_	_	-			-		_	2475
				gga Gly											2523
				cag Gln 800											2571
				cat His											2619
				ata Ile											2667

agc Ser	att Ile	cat His 845	cct Pro	cgg Arg	tca Ser	ttg Leu	tta Leu 850	ctg Leu	acg Thr	gga Gly	cca Pro	aac Asn 855	atg Met	ggc Gly	gga Gly	2715
														gcc Ala		2763
														gtg Val		2811
														gga Gly 905		2859
_			_	_	_	-				-				cag Gln		2907
														gga Gly		2955
														cac His		3003
														cac His		3051
		_	_							Val				cac His 985		3099
-	_	-						_				Arg		tgt Cys		3147
	Asp		Val			Tyr	_	Leu			Gly		Cys	cct Pro		3195
Ser		-			Val	_		_	_	Gly		Pro		caa Gln		3243
_	Glu		_	Ser		Ala	_		_		Lys				999 Gly 1050	3291

gga a																
Glu	aac Asn	ttc Phe	Lys	tca Ser 055	agt Ser	gag Glu	cta Leu	Arg	tct Ser 060	gag Glu	ttc Phe	tca Ser	Ser	ctg Leu .065	cat His	3339
gaa ( Glu		Trp					Val					Val				3387
aat (	Ala					Asp					Leu					3435
cat (					Ser						taaa	tggo	ta			3478
tgac	ataa	aca c	tato	tgaa	g ct	cgtt	aagt	ctt	ttgc	ctc	tctg	gatgt	tt a	attco	ctctta	3538
aaaa	atgo	ctt a	tata	tcaa	ıa aa	att	gttt	cto	gatt	aaa	aaaa	aaaa	aa a	aaaa	aaaaa	3598
aaaa	aaaa	a														3606
<210 <211 <212 <213 <223	>			: abido	-			na ec	otyp:	e Co	olumb	oia				
			POI	.yper	clae	MSI	16									
<400			31	.yper	cide	e MSI	16									
	>	Arg	31					Ser	Phe 10	Phe	Gln	Lys	Pro	Thr 15	Ala	
Met 1	> Gln	_	31 Gln	Arg 5	Ser	Ile	Leu		10			Lys Ser		15		
Met 1 Ala	o Gln Thr	Thr	31 Gln Lys 20	Arg 5 Gly	Ser Leu	Ile Val	Leu	Gly 25	10 Asp	Ala	Ala	-	Gly 30	15 Gly	Gly	
Met 1 Ala Gly	Gln Thr	Thr Gly 35	31 Gln Lys 20 Gly	Arg 5 Gly Pro	Ser Leu Arg	Ile Val Phe	Leu Ser Asn 40	Gly 25 Val	10 Asp Arg	Ala Glu	Ala	Ser	Gly 30	15 Gly Lys	Gly Gly	
Met 1 Ala Gly Asp	Gln Thr Ser Ala	Thr Gly 35 Ser	31 Gln Lys 20 Gly Val	Arg 5 Gly Pro	Ser Leu Arg	Ile Val Phe Ala 55	Leu Ser Asn 40 Val	Gly 25 Val Ser	10 Asp Arg Lys	Ala Glu Ser	Ala Gly Val 60	Ser Asp 45	Gly 30 Ala Glu	15 Gly Lys Val	Gly Gly Arg	
Met 1 Ala Gly Asp Gly 65	Gln Thr Ser Ala 50 Thr	Thr Gly 35 Ser Asp	31 Gln Lys 20 Gly Val	Arg 5 Gly Pro Arg	Ser Leu Arg Phe Pro 70	Ile Val Phe Ala 55 Glu	Leu Ser Asn 40 Val	Gly 25 Val Ser	10 Asp Arg Lys	Ala Glu Ser Arg 75	Ala Gly Val 60 Arg	Ser Asp 45	Gly 30 Ala Glu Leu	15 Gly Lys Val	Gly Gly Arg Ser 80	
Met 1 Ala Gly Asp Gly 65	Gln Thr Ser Ala 50 Thr	Thr Gly 35 Ser Asp	31 Gln Lys 20 Gly Val Thr	Arg 5 Gly Pro Arg Pro Ala 85	Ser Leu Arg Phe Pro 70 Glu	Ile Val Phe Ala 55 Glu Ser	Leu Ser Asn 40 Val Lys	Gly 25 Val Ser Val	Asp Arg Lys Pro Asp	Ala Glu Ser Arg 75 Ala	Ala Gly Val 60 Arg	Ser Asp 45 Asp	Gly 30 Ala Glu Leu	15 Gly Lys Val Pro	Gly Gly Arg Ser 80 Ser	

Met	Lys 130	Ala	Asn	Asp	Val	Ile 135	Pro	Gln	Phe	Arg	Ser 140	Asn	Asn	Gly	Lys
Thr 145	Gln	Glu	Arg	Asn	His 150	Ala	Phe	Ser	Phe	Ser 155	Gly	Arg	Ala	Glu	Leu 160
Arg	Ser	Val	Glu	Asp 165	Ile	Gly	Val	Asp	Gly 170	Asp	Val	Pro	Gly	Pro 175	Glu
Thr	Pro	Gly	Met 180	Arg	Pro	Arg	Ala	Ser 185	Arg	Leu	Lys	Arg	Val 190	Leu	Glu
Asp	Glu	Met 195	Thr	Phe	Lys	Glu	Asp 200	Lys	Val	Pro	Val	Leu 205	Asp	Ser	Asn
Lys	Arg 210	Leu	Lys	Met	Leu	Gln 215	Asp	Pro	Val	Cys	Gly 220	Glu	Lys	Lys	Glu
Val 225	Asn	Glu	Gly	Thr	Lys 230	Phe	Glu	Trp	Leu	Glu 235	Ser	Ser	Arg	Ile	Arg 240
Asp	Ala	Asn	Arg	Arg 245	Arg	Pro	Asp	Asp	Pro 250	Leu	Tyr	Asp	Arg	Lys 255	Thr
Leu	His	Ile	Pro 260	Pro	Asp	Val	Phe	Lys 265	Lys	Met	Ser	Ala	Ser 270	Gln	Lys
Gln	Tyr	Trp 275	Ser	Val	Lys	Ser	Glu 280	Tyr	Met	Asp	Ile	Val 285	Leu	Phe	Phe
Lys	Val 290	Gly	Lys	Phe	Tyr	Glu 295	Leu	Tyr	Glu	Leu	Asp 300	Ala	Glu	Leu	Gly
His 305	Lys	Glu	Leu	Asp	Trp 310	Lys	Met	Thr	Met	Ser 315	Gly	Val	Gly	Lys	Cys 320
Arg	Gln	Val	Gly	Ile 325	Ser	Glu	Ser	Gly	Ile 330		Glu	Ala	Val	Gln 335	Lys
Leu	Leu	Ala	Arg 340	_	Tyr	Lys	Val	Gly 345		, Ile	Glu	Gln	Leu 350	Glu	Thr
Ser	Asp	Gln 355		Lys	Ala	Arg	Gly 360		. Asn	Thr	Ile	Ile 365	Pro	Arg	Lys
Leu	Val 370		Val	Leu	Thr	Pro 375		Thr	Ala	. Ser	Glu 380		Asn	Ile	Gly
Pro 385	_	Ala	. Val	His	Leu 390		Ala	Ile	Lys	395		Lys	Met	Glu	Leu 400
Gln	Lys	Cys	Ser	Thr		Tyr	Gly	Phe	Ala 410		. Val	Asp	Cys	Ala 415	

Leu Arg Phe Trp Val Gly Ser Ile Ser Asp Asp Ala Ser Cys Ala Ala Leu Gly Ala Leu Leu Met Gln Val Ser Pro Lys Glu Val Leu Tyr Asp Ser Lys Gly Leu Ser Arg Glu Ala Gln Lys Ala Leu Arg Lys Tyr Thr Leu Thr Gly Ser Thr Ala Val Gln Leu Ala Pro Val Pro Gln Val Met Gly Asp Thr Asp Ala Ala Gly Val Arg Asn Ile Ile Glu Ser Asn Gly Tyr Phe Lys Gly Ser Ser Glu Ser Trp Asn Cys Ala Val Asp Gly Leu Asn Glu Cys Asp Val Ala Leu Ser Ala Leu Gly Glu Leu Ile Asn His Leu Ser Arg Leu Lys Leu Glu Asp Val Leu Lys His Gly Asp Ile Phe Pro Tyr Gln Val Tyr Arg Gly Cys Leu Arg Ile Asp Gly Gln Thr Met Val Asn Leu Glu Ile Phe Asn Asn Ser Cys Asp Gly Gly Pro Ser Gly Thr Leu Tyr Lys Tyr Leu Asp Asn Cys Val Ser Pro Thr Gly Lys Arg Leu Leu Arg Asn Trp Ile Cys His Pro Leu Lys Asp Val Glu Ser Ile Asn Lys Arg Leu Asp Val Val Glu Glu Phe Thr Ala Asn Ser Glu Ser Met Gln Ile Thr Gly Gln Tyr Leu His Lys Leu Pro Asp Leu Glu Arg Leu Leu Gly Arg Ile Lys Ser Ser Val Arg Ser Ser Ala Ser Val Leu Pro Ala Leu Leu Gly Lys Lys Val Leu Lys Gln Arg Val Lys Ala Phe Gly Gln Ile Val Lys Gly Phe Arg Ser Gly Ile Asp Leu Leu Leu Ala Leu Gln Lys Glu Ser Asn Met Met Ser Leu Leu Tyr Lys Leu Cys Lys

Leu 705	Pro	Ile	Leu	Val	Gly 710	Lys	Ser	Gly	Leu	Glu 715	Leu	Phe	Leu	Ser	720
Phe	Glu	Ala	Ala	Ile 725	Asp	Ser	Asp	Phe	Pro 730	Asn	Tyr	Gln	Asn	Gln 735	Asp
Val	Thr	Asp	Glu 740	Asn	Ala	Glu	Thr	Leu 745	Thr	Ile	Leu	Ile	Glu 750	Leu	Phe
Ile	Glu	Arg 755	Ala	Thr	Gln	Trp	Ser 760	Glu	Val	Ile	His	Thr 765	Ile	Ser	Cys
Leu	Asp 770	Val	Leu	Arg	Ser	Phe 775	Ala	Ile	Ala	Ala	Ser 780	Leu	Ser	Ala	Gly
Ser 785	Met	Ala	Arg	Pro	Val 790	Ile	Phe	Pro	Glu	Ser 795	Glu	Ala	Thr	Asp	Gln 800
Asn	Gln	Lys	Thr	Lys 805	Gly	Pro	Ile	Leu	Lys 810	Ile	Gln	Gly	Leu	Trp 815	His
Pro	Phe	Ala	Val 820	Ala	Ala	Asp	Gly	Gln 825	Leu	Pro	Val	Pro	Asn 830	Asp	Ile
Leu	Leu	Gly 835	Glu	Ala	Arg	Arg	Ser 840	Ser	Gly	Ser	Ile	His 845	Pro	Arg	Ser
Leu	Leu 850	Leu	Thr	Gly	Pro	Asn 855	Met	Gly	Gly	Lys	Ser 860	Thr	Leu	Leu	Arg
Ala 865	Thr	Cys	Leu	Ala	Val 870	Ile	Phe	Ala	Gln	Leu 875	Gly	Cys	Tyr	Val	Pro 880
Cys	Glu	Ser	Cys	Glu 885	Ile	Ser	Leu	Val	Asp 890	Thr	Ile	Phe	Thr	Arg 895	Leu
Gly	Ala	Ser	Asp 900	Arg	Ile	Met	Thr	Gly 905	Glu	Ser	Thr	Phe	Leu 910	Val	Glu
Cys	Thr	Glu 915	Thr	Ala	Ser	Val	Leu 920	Gln	Asn	Ala	Thr	Gln 925	Asp	Ser	Leu
Val	Ile 930	Leu	Asp	Glu	Leu	Gly 935	Arg	Gly	Thr	Ser	Thr 940	Phe	Asp	Gly	Tyr
Ala 945	Ile	Ala	Tyr	Ser	Val 950	Phe	Arg	His	Leu	Val 955	Glu	Lys	Val	Gln	Cys 960
Arg	Met	Leu	Phe	Ala 965	Thr	His	Tyr	His	Pro 970		Thr	Lys	Glu	Phe 975	Ala
Ser	His	Pro	Arg 980	Val	Thr	Ser	Lys	His 985	Met	Ala	Cys	Ala	Phe 990	Lys	Ser

acataaccac aaataggggt gc

Arg Ser Asp T	Tyr Gln Pro Arg Gly Cys Asp Gln Asp Leu Val Phe Leu 1000 1005	
Tyr Arg Leu T	Chr Glu Gly Ala Cys Pro Glu Ser Tyr Gly Leu Gln Val 1015 1020	
Ala Leu Met A 1025	Ala Gly Ile Pro Asn Gln Val Val Glu Thr Ala Ser Gly 1030 1035 1040	
Ala Ala Gln A	Ala Met Lys Arg Ser Ile Gly Glu Asn Phe Lys Ser Ser 1045 1050 1055	
_	Ser Glu Phe Ser Ser Leu His Glu Asp Trp Leu Lys Ser 1060 1065 1070	
Leu Val Gly 1	Ile Ser Arg Val Ala His Asn Asn Ala Pro Ile Gly Glu 1080 1085	
Asp Asp Tyr A	Asp Thr Leu Phe Cys Leu Trp His Glu Ile Lys Ser Ser 1095 1100	
Tyr Cys Val	Pro Lys	
<210>	32	
<211>	24	
<212> <213>	DNA Artificial sequence	
	-	
<220>		
<223>	Forward primer for PCR amplification of ATHGENEA microsatellite	
<400>	32	
accatgcata g	cttaaactt cttg	24
<210>	33	
<211>	22	
<212>	DNA	
<213>	Artificial sequence	
<220>	-	
<223>	Reverse primer for PCR amplification of ATHGENEA microsatellite	
<400>	33	

gctctagatt tctcgctcta agactctct

<210>	34
<211>	18
<212>	DNA
<213>	Artificial sequence
(213)	Aftificial sequence
<220>	
<223>	Forward primer DMCIN-A for PCR on genomic DNA of Arabidopsis
	thaliana ssp. Landsberg erecta "Ler"
<400>	34
gaagcgatat	tgttcgtg
<210>	35
<211>	18
<212>	DNA
<213>	Artificial sequence
<220>	•
	December 2017 D. Son DCD on account DNA of Auchidensi
<223>	Reverse primer DMCIN-B for PCR on genomic DNA of Arabidopsi thaliana ssp. Landsberg erecta "Ler"
<400>	35
agattgcgag	aacattcc 1
-210-	36
<210>	
<211>	31
<212>	DNA
<213>	Artificial sequence
<220>	
<223>	Forward primer DMCIN-1 for PCR on genomic DNA of Arabidopsi
	thaliana ssp. Landsberg erecta "Ler"
<400>	36
acgcgtcgac	tcagctatga gattactcgt g 3
<210>	27
<210>	37
	29
<212>	DNA
<213>	Artificial sequence
<220>	$oldsymbol{\epsilon}$
<223>	Reverse primer DMCIN-2 for PCR on genomic DNA of Arabidopsi
	thaliana ssp. Landsberg erecta "Ler"
<400>	37

<210>	38
<211>	32
<212>	DNA
<213>	Artificial sequence
<220>	The state of the s
<223>	Forward primer DMCIN-3 for PCR on genomic DNA of Arabidopsis thaliana ssp. Landsberg erecta "Ler"
<400>	38
gctctagagc	ttctcttaag taagtgattg at 32
210	
<210>	39
<211>	48
<212>	DNA
<213>	Artificial sequence
<220>	
<223>	Reverse primer DMCIN-4 for PCR on genomic DNA of Arabidopsis
	thaliana ssp. Landsberg erecta "Ler"
<400>	39
tecceeggge	togagagato tocatggttt ottoagetot atgaatce 48
<210>	40
<211>	26
<212>	DNA
<213>	Artificial sequence
<220>	
<223>	Forward primer DMCla for PCR on genomic DNA of Arabidopsis
	thaliana ssp. Landsberg erecta "Ler"
<400>	40
acgcgtcgac	gaattcgcaa gtgggg 26
<210>	41
<211>	38
<212>	DNA
<213>	Artificial sequence
<220>	
<223>	Reverse primer DMC1b for PCR on genomic DNA of Arabidopsis
~667/	thaliana ssp. Landsberg erecta "Ler"
<400>	41

tccatggaga	totocogggt accgattige ttogaggg	38
33 3	3 333	
<210>	42	
<211>	20	
<212>	DNA	
<213>	Artificial sequence	
	merrical sequence	
<220>	•	
<223>	Forward primer for PCR amplification of ATEAT1 SSLP marker	in
	Arabidopsis thaliana subspecies	
<400>	42	
accactacat	gaatgatatg	20
gccactgcgt	gaacgacacg	20
<210>	43	
<211>	22	
<212>	DNA	
<213>	Artificial sequence	
<220>		
<223>	Reverse primer for PCR amplification of ATEAT1 SSLP marker	· in
	Arabidopsis thaliana subspecies	
<400>	43	
cgaacagcca	acattaattc cc	22
<210>	44	
<211>	18	
<212>	DNA	
<213>	Artificial sequence	
<220>		
<223>	Forward primer for PCR amplification of NGA63 SSLP marker	in
	Arabidopsis thaliana subspecies	
<400>	44	
aaccaaggca	Cagaagcg	18
		10
<210>	45	
<211>	18	
<211>		
	DNA	
<213>	Artificial sequence	
<220>		
<223>	Reverse primer for PCR amplification of NGA63 SSLP marker Arabidopsis thaliana subspecies	in

<400>	45	
acccaagtga	tcgccacc	18
<210>	46	
<211>	21	
<212>	DNA	
<213>	Artificial sequence	
<220>		
<223>	Forward primer for PCR amplification of NGA248 SSLP m Arabidopsis thaliana subspecies	arker in
<400>	46	
taccgaacca	aaacacaaag g	21
<210>	47	
<211>	22	
<212>	DNA	
<213>	Artificial sequence	
(213)	Artificial bodacioc	
<220>		
<223>	Reverse primer for PCR amplification of NGA248 SSLP m Arabidopsis thaliana subspecies	arker in
<400>	47	
tctgtatctc	ggtgaattet ee	22
<210>	48	
<211>	22	
<212>	DNA	
<213>	Artificial sequence	
<220>		
<223>	Forward primer for PCR amplification of NGA128 SSLP m Arabidopsis thaliana subspecies	arker in
<400>	48	
ggtctgttga	tgtcgtaagt cg	22
<210>	49	
<211>	22	
<211>	DNA	
<212>	Artificial sequence	
~~~/	WICITIOIAL Seducace	
<220>		

PCT/EP98/06977 -WO 99/19492

	37	• • • • • • • • • • • • • • • • • • • •
<223>	Reverse primer for PCR amplification of NGA128 SSLP Arabidopsis thaliana subspecies	marker in
<400>	4 9	
atcttgaaac	ctttagggag gg	22
<210>	50	
<211>	22	
<212>	DNA	
<213>	Artificial sequence	
<220>		
<223>	Forward primer for PCR amplification of NGA280 SSLP Arabidopsis thaliana subspecies	marker in
<400>	50	
ctgatctcac	ggacaatagt gc	22
<210>	51	
<211>	20	
<212>	DNA	
<213>	Artificial sequence	
<220>		
<223>	Reverse primer for PCR amplification of NGA280 SSLP Arabidopsis thaliana subspecies	marker in
<400>	51	
ggctccataa	aaagtgcacc	20
<210>	52	
<211>	21	
<212>	DNA	
<213>	Artificial sequence	
<220>		
<223>	Forward primer for PCR amplification of NGA111 SSLP Arabidopsis thaliana subspecies	marker in
<400>	52	
ctccagttgg	aagctaaagg g	21
<210>	53	
<211>	21	
<212>	DNA	

<213> Artificial sequence

PCT/EP98/06977 WO 99/19492

	38	
<220>		
<223>	Reverse primer for PCR amplification of NGA111 SSLP market Arabidopsis thaliana subspecies	r in
<400>	53	
tgttttttag g	gacaaatggc g	21
<210>	54	
<211>	20	
<212>	DNA	
<213>	Artificial sequence	
<220>		
<223>	Forward primer for PCR amplification of NGA168 SSLP marker Arabidopsis thaliana subspecies	: ir
<400>	54	
ccttcacatc o	caaaacccac	2.0
<210>		
<211>	55 20	
<211>		
<213>	DNA	
<213>	Artificial sequence	
<220>		
<223>	Reverse primer for PCR amplification of NGA168 SSLP marker Arabidopsis thaliana subspecies	: ir
<400>	55	
gcacataccc a	acaaccagaa	20
<210>	56	
<211>	20	
<212>	DNA	
<213>	Artificial sequence	
<220>		
<223>	Forward primer for PCR amplification of NGA1126 SSLP marke in Arabidopsis thaliana subspecies	r

cgctacgctt ttcggtaaag

56

<400>

PCT/EP98/06977 WO 99/19492

39

<210>	57
<211>	20
<212>	DNA
<213>	Artificial sequence
(213)	Artificial sequence
<220>	
<223>	Reverse primer for PCR amplification of NGA1126 SSLP marker in Arabidopsis thaliana subspecies
<400>	57
gcacagtcca a	gtcacaacc 20
<210>	58
<211>	20
<212>	DNA
<213>	Artificial sequence
<220>	
<223>	Forward primer for PCR amplification of NGA361 SSLP marker in Arabidopsis thaliana subspecies
<400>	58
aaagagatga g	aatttggac 20
<210>	59
<211>	23
<212>	DNA
<213>	Artificial sequence
<220>	
<223>	Reverse primer for PCR amplification of NGA361 SSLP marker in Arabidopsis thaliana subspecies
<400>	59
acatatcaat a	tattaaagt agc 23
<210>	60
<211>	18
<212>	DNA
<213>	
~213/	Artificial sequence
<220>	
<223>	Forward primer for PCR amplification of NGA168 SSLP marker in Arabidopsis thaliana subspecies

<400> 60

tegtetactg	Cactgccg 18
<210>	61
<211>	22
<212>	DNA
<213>	Artificial sequence
<220>	
<223>	Reverse primer for PCR amplification of NGA168 SSLP marker in Arabidopsis thaliana subspecies
<400>	61
gaggacatgt	ataggageet eg 22
<210>	62
<211>	20
<212>	DNA
<213>	Artificial sequence
<220>	
<223>	Forward primer for PCR amplification of AthBIO2 SSLP marker
	in Arabidopsis thaliana subspecies
<400>	62
tgacctcctc	ttccatggag 20
<210>	63
<211>	22
<212>	DNA
<213>	Artificial sequence
<220>	
<223>	Reverse primer for PCR amplification of AthBIO2 SSLP marker
	in Arabidopsis thaliana subspecies
<400>	63
ttaacagaaa	cccaaagett te 22
<210>	64
<211>	21
<212>	DNA
<213>	Artificial sequence
~~~~	ererreda pedreuce
<220>	
<223>	Forward primer for PCR amplification of AthUBIQUE SSLP marker in Arabidopsis thaliana subspecies

<400>	64
aggcaaatgt	ccatttcatt g 21
<210>	65
<211>	20
<212>	DNA
<213>	Artificial sequence
<220>	
<223>	Reverse primer for PCR amplification of AthUBIQUE SSLP marker in Arabidopsis thaliana subspecies
<400>	65
acgacatggc	agatttctcc 20
<210>	66
<211>	
	21
<212>	DNA
<213>	Artificial sequence
<220>	
<223>	Forward primer for PCR amplification of NGA172 SSLP marker in Arabidopsis thaliana subspecies
<400>	66
agctgcttcc	ttatagcgtc c 21
<210>	
<210>	67
<211>	19
	DNA
<213>	Artificial sequence
<220>	
<223>	Reverse primer for PCR amplification of NGA172 SSLP marker in Arabidopsis thaliana subspecies
<400>	67
catccgaatg	ccattgttc 19
<210>	60
	68
<211>	21
<212>	DNA
<213>	Artificial sequence
<220>	

Artificial sequence

<213>

	42					
<223>	Forward primer for PCR amplification Arabidopsis thaliana subspecies	of	NGA126	SSLP	marker	in
<400>	68					
gaaaaaacgc	tactttcgtg g					21
<210>	69					
<211>	22					
<212>	DNA					
<213>	Artificial sequence					
<220>						
<223>	Reverse primer for PCR amplification Arabidopsis thaliana subspecies	of	NGA126	SSLP	marker	in
<400>	69					
caagagcaat	atcaagagca gc					22
<210>	70					
<211>	20					
<212>	DNA					
<213>	Artificial sequence					
<220>	•					
<223>	Forward primer for PCR amplification Arabidopsis thaliana subspecies	of	NGA162	SSLP	marker	ir
<400>	70					
catgcaattt	gcatctgagg					20
<210>	71					
<211>	22					
<212>	DNA					
<213>	Artificial sequence					
<220>						
<223>	Reverse primer for PCR amplification Arabidopsis thaliana subspecies	of	NGA162	SSLP	marker	in
<400>	71					
ctctgtcact	cttttcctct gg					22
<210>	72					
<211>	21					
<212>	DNA					
	₩4143					

PCT/EP98/06977

<220>		
<223>	Forward primer for PCR amplification of NGA6 SSLP marker in Arabidopsis thaliana subspecies	.n
<400>	72	
tggatttctt	cctctcttca c	21
<210>	73	
<211>	21	
<212>	DNA	
<213>	Artificial sequence	
<220>		
<223>	Reverse primer for PCR amplification of NGA6 SSLP marker i Arabidopsis thaliana subspecies	n.
<400>	73	
atggagaagc	ttacactgat c	21
<210>	74	
<211>	20	
<212>	DNA	
<213>	Artificial sequence	
(213)	Altilitai sequence	
<220>		
<223>	Forward primer for PCR amplification of NGA12 SSLP marker Arabidopsis thaliana subspecies	in
<400>	74	
aatgttgtcc	tcccctcctc	20
<210>	75	
<211>	22	
<212>	DNA	
<213>	Artificial sequence	
<220>		
<223>	Reverse primer for PCR amplification of NGA12 SSLP marker	in
- M 40 07 F	Arabidopsis thaliana subspecies	T #1
<400>	75	
tgatgctctc	tgaaacaaga gc	22

WO 99/19492

44

<210>	
<211>	76 21
<212>	DNA
<213>	Artificial sequence
,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,	Artiritat sequence
<220>	
<223>	Forward primer for PCR amplification of NGA8 SSLP marker in Arabidopsis thaliana subspecies
<400>	76
gagggcaaa	t ctttatttcg g
<210>	77
<211>	22
<212>	DNA
<213>	Artificial sequence
<220>	
<223>	Reverse primer for PCR amplification of NGA8 SSLP marker in
	Arabidopsis thaliana subspecies
<400>	77
tggctttcg	t ttataaacat cc 22
<210>	78
<211>	21
<212>	DNA
<213>	Artificial sequence
<220>	
<223>	Forward primer for PCR amplification of NGA1107 SSLP marker in Arabidopsis thaliana subspecies
<400>	78
gcgaaaaaa	c aaaaaaatcc a 21
1	
<210>	79
<211>	21
<212>	DNA
<213>	Artificial sequence
<220>	
<223>	Reverse primer for PCR amplification of NGA1107 SSLP marker in Arabidopsis thaliana subspecies
<400>	79

cgacgaatcg	acagaattag g	21
<210>	80	
<211>	21	
<212>	DNA	
<213>	Artificial sequence	
<220>		
<223>	Forward primer for PCR amplification of NGA225 SSLP marker Arabidopsis thaliana subspecies	in
<400>	80	
gaaatccaaa	tcccagagag g	21
-210-		
<210>	81	
<211>	22	
<212>	DNA	
<213>	Artificial sequence	
<220>		
<223>	Reverse primer for PCR amplification of NGA225 SSLP marker Arabidopsis thaliana subspecies	in
<400>	81	
tctccccact	agttttgtgt cc	22
<210>	82	
<211>	19	
<212>	DNA	
<213>	Artificial sequence	
<220>		
<223>	Forward primer for PCR amplification of NGA249 SSLP marker Arabidopsis thaliana subspecies	in
<400>	82	
taccgtcaat	ttcatcgcc	١9
<210>	83	
<211>	22	
<212>	DNA	
<213>	Artificial sequence	
<220>		
<223>	Reverse primer for PCR amplification of NGA249 SSLP marker	in
	Arabidopsis thaliana subspecies	T- 1.1

46

<400>	83
ggatosotaa	ctgtaaaatc cc 22
<210>	84
<211>	22
<212>	DNA
<213>	Artificial sequence
<220>	
<223>	Forward primer for PCR amplification of CA72 SSLP marker in Arabidopsis thaliana subspecies
<400>	84
aatcccagta a	accaaacaca ca 22
<210>	85
<211>	20
<212>	
	DNA Nami Giania I
<213>	Artificial sequence
<220>	
<223>	Reverse primer for PCR amplification of CA72 SSLP marker in
	Arabidopsis thaliana subspecies
<400>	85
cccagtctaa d	ccacgaccac 20
<210>	86
<211>	20
<212>	DNA
<213>	Artificial sequence
	ALCILICIAI Sequence
<220>	
<223>	Forward primer for PCR amplification of NGA151 SSLP marker in Arabidopsis thaliana subspecies
<400>	86
gttttgggaa g	gttttgctgg 20
<210>	87
<211>	24
<212>	DNA
<213>	Artificial sequence
<220>	

<223>	Reverse primer for PCR amplification of NGA151 SSLP marker Arabidopsis thaliana subspecies	in
<400>	87	
cagtctaaaa	gcgagagtat gatg	24
<210>	,88 <sup></sup>	
<211>	22	
<212>	DNA	
<213>	Artificial sequence	
(213)	Artificial Sequence	
<220>		
<223>	Forward primer for PCR amplification of NGA106 SSLP marker Arabidopsis thaliana subspecies	in
<400>	88	
gttatggagt	ttctagggca cg	22
<210>	8 9	
<211>	20	
<212>	DNA	
<213>	Artificial sequence	
<220>		
<223>	Reverse primer for PCR amplification of NGA106 SSLP marker	i n
	Arabidopsis thaliana subspecies	111
<400>	8 9	
tgccccattt	tgttcttctc	20
<210>	90	
<211>	20	
<212>	DNA	
<213>	Artificial sequence	
<220>		
<223>	Forward primer for PCR amplification of NGA139 SSLP marker	in
	Arabidopsis thaliana subspecies	111
<400>	90	
agagctacca	gatecgatgg	20
<210>	91	
<211>	21	
<212>	DNA	
<213>	Artificial sequence	

PCT/EP98/06977

WO 99/19492 PCT/EP9

<220> <223>	Reverse primer for PCR amplification of NGA139 SSLP marker in Arabidopsis thaliana subspecies
<400>	91
ggtttcgttt	cactatecag g 21
<210><211><212><213>	92 22 DNA Artificial sequence
<220> <223>	Forward primer for PCR amplification of NGA76 SSLP marker in Arabidopsis thaliana subspecies
<400>	92
ggagaaaatg	tcactctcca cc 22
<210> <211> <212> <213> <220> <223>	93 20 DNA Artificial sequence  Reverse primer for PCR amplification of NGA76 SSLP marker in
(223)	Arabidopsis thaliana subspecies
<400>	93
aggcatggga	gacatttacg 20
<210> <211> <212> <213>	94 20 DNA Artificial sequence
<220> <223>	Forward primer for PCR amplification of ATHSO191 SSLP marker in Arabidopsis thaliana subspecies
<400>	94

ctccaccaat catgcaaatg

20

49

<210>	95
<211>	21
<212>	DNA
<213>	Artificial sequence
<220>	
<223>	Reverse primer for PCR amplification of ATHSO191 SSLP marker in Arabidopsis thaliana subspecies
<400>	95
tgatgttgat g	gagatggtc a 21
<210>	96
<211>	22
<212>	DNA
<213>	Artificial sequence
<220>	
<223>	Forward primer for PCR amplification of NGA129 SSLP marker in
	Arabidopsis thaliana subspecies
<400>	96
tcaggaggaa c	taaagtgag gg 22
<210>	97
<211>	22
<212>	DNA
<213>	Artificial sequence
<220>	
<223>	Reverse primer for PCR amplification of NGA129 SSLP marker in Arabidopsis thaliana subspecies
<400>	97
cacactgaag a	tggtcttga gg 22
<210>	98
<211>	8062
<212>	DNA
<213>	Arabidopsis thaliana ecotype Columbia
<220>	
<223>	Genomic DNA sequence of AtMSH6
<400>	97

aagaaatgaa	agatatatat	tgttttttca	tttatcaaac	aaaacaacaa	gactttttt	120
ttacttttta	cattggtcaa	Caaaatacaa	gataaacgac	atcgtttaat	catttcccaa	180
tttacccct	aagtttaaca	cctagaacct	tctccatctt	cgcaagcaca	gcctgattag	240
gaacagcttt	accattctca	tattcctgaa	ctacctgagt	cctctcattg	atctgtttcg	300
ccaaatccgc	ttgtgacatc	ttcttctcca	atctcgcttt	ctgtatcatc	aacctcacct	360
ctgctttcac	acgatccatc	gccgcaggct	ctgtttcttc	ttccagcttc	ttcgtgttaa	420
tcaccggaac	cgccgtagat	ttcccctttt	tgttcgaacc	ggcatcgaat	ttcttaaccg	480
tttgaaccgc	gacaccgttt	ctcagagctg	cgttaaccgc	tttcggatcg	cgtaggtctt	540
ggctcttttg	ttttgatttg	tggagaacta	ctggttccca	gtcttgtgtt	actgctcctg	600
ggtatctgct	cggcatcgtc	gatgaattga	gagaaaggaa	caacgcgaaa	attttattaa	660
tctgagtttt	gaaattgaga	aacgatgaag	atgaagaatg	ttgttgagag	gattgtgata	720
tttatatata	cgaagattgg	tttctggaga	attcgatcat	ctttttatca	attttcgtct	780
ctggaacgtt	cttagagatg	attgacgacg	tgtcattatc	tgatttgcag	ttaaccaatg	840
ctttttgggt	tggattcgtg	gtacaccata	ttatccgatt	tggctcaatg	gttttatata	900
aatttggttt	teggtteggt	tatgagttat	cattaaaatt	aagctaacca	aaaattttcg	960
taaaatttat	ttcggtttca	attcggatcc	cttacttcca	gaaccgaatt	attcgaaacc	1020
ggggttagcc	gaaccgaata	ccaatgcctg	attgactcgt	tggctagaaa	gatccaacgg	1080
tatacaataa	tagaacataa	atcggacggt	catcaaagcc	tcaaagagtg	aacagtcaac	1140
aaaaaagtt	gagecetgag	gagtatcgtt	tccgccattt	ctacgacgca	aggcgaaaat	1200
ttttggcgcc	aatctttccc	ccctttcgaa	ttctctcagc	tcaaaacatc	gtttctctct	1260
cactctctct	cacaattcca	aaaaatgcag	cgccagagat	cgattttgtc	tttcttccaa	1320
aaacccacgg	cggcgactac	gaagggtttg	gtttccggcg	atgctgctag	cggcggggc	1380
ggcagcggag	accacgattt	aatgtgaagg	aaggggatgc	taaaggcgac	gcttctgtac	1440
gttttgctgt	ttcgaaatct	gtcgatgagg	ttagaggaac	ggatactcca	ccggagaagg	1500
ttccgcgtcg	tgtcctgccg	tctggattta	agccggctga	atccgccggt	gatgcttcgt	1560
ccctgttctc	caatattatg	cataagtttg	taaaagtcga	tgatcgagat	tgttctggag	1620
agaggtacta	atcttcgatt	ctcttaattt	tgttatcttt	agctggaaga	agaagattcg	1680

tgtaatttgt	tgtattcgtt	ggagagattc	tgattactgc	attggatcgt	tgtttacaaa	1740
ttttcaggag	ccgagaagat	gttgttccgc	tgaatgattc	atctctatgt	atgaaggcta	1800
atgatgttat	tcctcaattt	cgttccaata	atggtaaaac	tcaagaaaga	aaccatgctt	1860
ttagtttcag	tgggagagct	gaacttagat	cagtagaaga	tataggagta	gatggcgatg	1920
ttcctggtcc	agaaacacca	gggatgcgtc	cacgtgcttc	tcgcttgaag	cgagttctgg	1980
aggatgaaat	gacttttaag	gaggataagg	ttcctgtatt	ggactctaac	aaaaggctga	2040
aaatgctcca	ggatccggtt	tgtggagaga	agaaagaagt	aaacgaagga	accaaatttg	2100
aatggcttga	gtcttctcga	atcagggatg	ccaatagaag	acgtectgat	gateceettt	2160
acgatagaaa	gaccttacac	ataccacctg	atgttttcaa	gaaaatgtct	gcatcacaaa	2220
agcaatattg	gagtgttaag	agtgaatata	tggacattgt	gcttttcttt	aaagtggtta	2280
gtaactatta	atctagtgtt	caatccattt	cctcaatgtg	atttgttcac	ttacatctgt	2340
ttacgttatg	ctcttctcag	gggaaatttt	atgagctgta	tgagctagat	gcggaattag	2400
gtcacaagga	gcttgactgg	aagatgacca	tgagtggtgt	gggaaaatgc	agacaggtaa	2460
attagttgaa	acaactggcc	tgcttgaatt	attgtgtcta	taaattttga	caccaccttt	2520
tgtttcaggt	tggtatctct	gaaagtggga	tagatgaggc	agtgcaaaag	ctattagctc	2580
gtgggtaagg	gaaccatcat	actttatgga	attcgtttac	tgctacttcg	gctaggattt	2640
aagaaatgga	aatcacttca	agcatcatta	gttaggatcc	tgagaactca	ggatgttttc	2700
ttattcgtta	tataataagt	cttttcatca	aggagtaaca	aacaaaactt	gcacaatatt	2760
tgtgtgctca	ctggcaaggc	atatataccc	agctaacctt	tgctagttca	ctgtagtaac	2820
agttacggat	aatatatgtt	tacttgtatg	tggtaccctc	attttgtctc	tcatggaggc	2880
tttcaagcct	tgtgttgaaa	ctggatagtt	acatatgctt	ccaacagaaa	ctagcatgca	2940
gattcatatg	ctttcctatt	ctactaatta	tgtattgaca	cactcgttgt	ttcttttgaa	3000
agatataaag	ttggacgaat	cgagcagcta	gaaacatctg	accaagcaaa	agccagaggt	3060
gctaatactg	taagttttct	tggataggtc	aaggagagtg	ttgcagactg	tttttgatca	3120
tttcttttc	tgtacattac	tttcatgctg	taattaactc	aatggctatt	ctggtctgat	3180
tatcagataa	ttccaaggaa	gctagttcag	gtattaactc	catcaacagc	aagcgaggga	3240
aacatcgggc	ctgatgccgt	ccatcttctt	gctataaaag	aggtttgtta	tttacttatt	3300

tatcttatca	tgttcagttc	atccaagtcc	tgaaaaatta	cactettett	taccaatctt	3360
ccatcaagct	gtgtaaagga	tttggaatta	gaaaatcatt	atttgatgct	ttgttttata	3420
tgcaagaggt	tcccttgaaa	agatctgttt	aagattcttt	gcacttgaaa	aattcaatct	3480
ttttaagtga	atcccctact	ttcttacaat	gatcatagtc	tgcaattgca	tgtcaagtaa	3540
tatcattcct	tgttactgca	teccetett	tcttaatgac	cattgtctat	gttgtgtttg	3600
tctcgtgtgc	tggagaaaat	gatagctgat	ccaagctgta	cattatcatg	attaagtagc	3660
tgctcaggaa	ttgcctttgg	ttacattgcc	taatggtttg	atgtcaattt	ttcttctgaa	3720
tctttatttt	agatcaaaat	ggagctacaa	aagtgttcaa	ctgtgtatgg	atttgctttt	3780
gttgactgtg	ctgccttgag	gttttgggtt	gggtccatca	gcgatgatgc	atcatgtgct	3840
gctcttggag	cgttattgat	gcaggtaagc	aagtgtattc	tgtatcttat	gtgtaccatg	3900
tgacttcctg	tgcatatatt	tgggttgcag	gaactaattc	tgaatcacca	tttggtatgt	3960
ttttccagg	tttctccaaa	ggaagtgtta	tatgacagta	aaggtaaact	gcttgtatcg	4020
ccagttgttt	tgttaaacag	aatttaaggt	aaatgacact	ggttaattta	aagtgcatac	4080
atgttgaaat	attgcagggc	tatcaagaga	agcacaaaag	gctctaagga	aatatacgťt	4140
gacaggtacc	atttcagtag	gcaagctaac	tgacaattta	accgctcacc	gaatgatagg	4200
tctcttaaac	attgctaatg	tagatgatgt	ttatgtttca	atctaatagg	gtctacggcg	4260
gtacagttgg	ctccagtacc	acaagtaatg	ggggatacag	atgctgctgg	agttagaaat	4320
ataatagaat	ctaacggata	ctttaaaggt	tcttctgaat	catggaactg	tgctgttgat	4380
ggtctaaatg	aatgtgatgt	tgcccttagt	gctcttggag	agctaattaa	tcatctgtct	4440
aggctaaagg	tgtgttggct	tgtttagttt	ttgcttttca	caaattaagc	aaaggaactt	4500
ttcataactt	acagtttcta	tctacttgca	gctagaagat	gtacttaagc	atggggatat	4560
ttttccatac	caagtttaca	ggggttgtct	cagaattgat	ggccagacga	tggtaaatct	4620
tgagatattt	aacaatagct	gtgatggtgg	tccttcaggc	aagtgcatat	ttcttttttg	4680
ataacttcaa	ctagagggca	gacatagaag	gaaaaattct	aatacttcgt	acggatctcc	4740
agtaagtaat	agccgatttt	tgtttaccta	tgtagggacc	ttgtacaaat	atcttgataa	4800
ctgtgttagt	ccaactggta	agcgactctt	aaggaattgg	atctgccatc	cactcaaaga	4860
tgtagaaagc	atcaataaac	ggcttgatgt	agttgaagaa	ttcacggcaa	actcagaaag	4920

tatgcaaatc	actggccagt	atctccacaa	acttccagac	ttagaaagac	tgctcggacg	4980
catcaagtct	agcgttcgat	catcagcctc	tgtgttgcct	gctcttctgg	ggaaaaaagt	5040
gctgaaacaa	cgagtaagta	tcaatcacaa	gttttctgag	taatgccttc	catgagtagt	5100
ataggactaa	aacattacgg	gtctagctaa	agactgttct	ccttcttttg	caatgtctgg	5160
ttattcatta	catttctctt	aacttattgc	attgcaggtt	aaagcatttg	ggcaaattgt	5220
gaaagggttc	agaagtggaa	ttgatctgtt	gttggctcta	cagaaggaat	caaatatgat	5280
gagtttgctt	tataaactct	gtaaacttcc	tatattagta	ggaaaaagcg	ggctagagtt	5340
atttctttct	caattcgaag	cagccataga	tagcgacttt	ccaaattatc	aggtgcccat	5400
ctatctttca	tactttacaa	caaaatgtct	gtcactactc	aaagcaatgc	atatggctta	5460
gatctcaact	cacaccccga	ggatcctaaa	gggatttgct	ttttattcct	aatgtttttg	5520
gatggtttga	tttatttcta	acttgaactt	attaatcttg	taccagaacc	aagatgtgac	5580
agatgaaaac	gctgaaactc	tcacaatact	tatcgaactt	tttatcgaaa	gagcaactca	5640
atggtctgag	gtcattcaca	ccataagctg	cctagatgtc	ctgagatctt	ttgcaatcgc	5700
agcaagtctc	tctgctggaa	gcatggccag	gcctgttatt	tttcccgaat	cagaagctac	5760
agatcagaat	cagaaaacaa	aagggccaat	acttaaaatc	caaggactat	ggcatccatt	5820
tgcagttgca	gccgatggtc	aattgcctgt	tccgaatgat	atactccttg	gcgaggctag	5880
aagaagcagt	ggcagcattc	atcctcggtc	attgttactg	acgggaccaa	acatgggcgg	5940
aaaatcaact	cttcttcgtg	caacatgtct	ggccgttatc	tttgcccaag	tttgtatact	6000
cgttagataa	ttactctatt	ctttgcaatc	agttcttcaa	catgaataat	aaattctgtt	6060
ttctgtctgc	agcttggctg	ctacgtgccg	tgtgagtctt	gcgaaatctc	cctcgtggat	6120
actatcttca	caaggcttgg	cgcatctgat	agaatcatga	caggagagag	taagttttgt	6180
tctcaaaata	ccaattcctc	gaactattta	ctcagatttt	gtctgattgg	acaaggtggt	6240
tttgcttttt	tttaggtacc	tttttggtag	aatgcactga	gacagcgtca	gttcttcaga	6300
atgcaactca	ggattcacta	gtaatccttg	acgaactggg	cagaggaact	agtactttcg	6360
atggatacgc	cattgcatac	tcggtaacct	gctcttctcc	ttcaacttat	acttgttgat	6420
caacaaaaac	atgcaattca	ttttgctgaa	acttattgat	ttatatcagg	tttttcgtca	6480
cctggtagag	aaagttcaat	gtcggatgct	ctttgcaaca	cattaccácc	ctctcaccaa	6540

ggaattcgcg	tctcacccac	gtgtcacctc	gaaacacatg	gcttgcgcat	tcaaatcaag	6600
atctgattat	caaccacgtg	gttgtgatca	agacctagtg	ttcttgtacc	gtttaaccga	6660
gggagcttgt	cctgagagct	acggacttca	agtggcactc	atggctggaa	taccaaacca	6720
agtggttgaa	acagcatcag	gtgctgctca	agccatgaag	agatcaattg	gggaaaactt	6780
caagtcaagt	gagctaagat	ctgagttctc	aagtctgcat	gaagactggc	tcaagtcatt	6840
ggtgggtatt	tctcgagtcg	cccacaacaa	tgccccatt	ggcgaagatg	actacgacac	6900
tttgttttgc	ttatggcatg	agatcaaatc	ctcttactgt	gttcccaaat	aaatggctat	6960
gacataacac	tatctgaagc	tcgttaagtc	ttttgcttct	ctgatgttta	ttcctcttaa	7.020
aaaatgctta	tatatcaaaa	aattgtttcc	tcgattataa	caagattata	tatgtatctg	7080
tcggtttagc	tatggtatat	aatatatgta	tgttcatgag	attggtcaag	agaaatactc	7140
acaaacagta	tattaagaag	gaaatatgtt	tatgcattaa	tttaagtttc	aagataaact	7200
gcaaataacc	tcgactaaag	ttgcaaagac	caaacacaaa	ttacaaaact	tataagactt	7260
aagttctgaa	ttccctaaaa	ccaaaaaaaa	aaacagaaca	tattttgttg	catctacaaa	7320
caacacaaac	ctacatagtt	tataacttac	tcatcactga	gattaacatc	agaatcattc	7380
tccatttctt	catcttcact	ctcatcatca	tcaccaccac	catgatgatt	ctcctcct	7440
tcacgtaacc	tagcaatctc	actctgagct	ctatcaacaa	tctgcttctt	ctgcaactcc	7500
aaatctctct	gaaaatcagc	tctcatcttc	tccaactcct	tcatttgctc	tttcttactc	7560
ttctccatct	tctcataaac	cttcccaaac	ctctcaacag	aatccgccaa	catcttatac	7620
gaagcagcgt	cattaacctt	cttcctctcg	tactcaacct	catcatcctc	atcctcctcc	7680
tcttcagaat	caccaggact	atccatcatc	tcatcaaacc	cattagactt	atctaaataa	7740
accttagtgt	tcataaacac	aaactcacct	gaatcaacac	cacaagctaa	acctaaatcc	7800
gacttgggcg	aaacacaaag	caacatatcc	aacttattga	aaaacgacca	tttacttgaa	7860
cctaaacctg	atttctcaac	cttaatcttc	tetttetat	acttcctctt	caagtcatca	7920
atcattctcc	tacattgcgt	ctcagatttc	tccatcctta	gctcctcact	cactttctca	7980
gctacttcat	tccaatcctc	gttcctcaaa	ctccttctac	ccaattgcaa	aaacctatct	8040
ccccaaactt	саадсаасас	aa				8062

## PCT

# WORLD INTELLECTUAL PROPERTY ORGANIZATION International Bureau



### INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification <sup>6</sup>: C12N 15/82, 15/29, C07K 14/415, C12N 15/10, 5/04, 5/14, A01H 5/00

A3 (11) International Publication Number:

WO 99/19492

(43) International Publication Date:

22 April 1999 (22.04.99)

(21) International Application Number:

PCT/EP98/06977

(22) International Filing Date:

9 October 1998 (09.10.98)

(30) Priority Data:

PO 9745

10 October 1997 (10.10.97) AU

(71) Applicant (for all designated States except US):
RHONE-POULENC AGRO [FR/FR]; 14/20, rue Pierre
Baizet, F-69009 Lyon (FR).

(72) Inventors; and

- (75) Inventors/Applicants (for US only): DOUTRIAUX,
  Marie-Pascale [FR/FR]; 64, route de Villebon, F-91160
  Saulx les Chartreux (FR). BETZNER, Andreas, Stefan
  [AU/AU]; 40 Dallachy Place, Page, ACT 2614 (AU).
  FREYSSINET, Georges [FR/FR]; 21, rue de Nervieux,
  F-69450 Saint Cyr au Mont d'Or (FR). PEREZ, Pascal
  [FR/FR]; 17, chemin de la Pradelle, Varennes, F-63450
  Chanonat (FR).
- (74) Agent: GENIN, Patrick; Rhône-Poulenc Agro, DPI, 14/20, rue Pierre Baizet, F-69009 Lyon (FR).

(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

#### Published

With international search report.

Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.

(88) Date of publication of the international search report:
24 June 1999 (24.06.99)

(54) Title: METHODS FOR OBTAINING PLANT VARIETIES

(57) Abstract

An isolated and purified DNA molecule comprising a polynucleotide sequence encoding a polypeptide functionally involved in the DNA mismatch repair system of a plant.

## FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
ΑU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ.	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	LT	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav	TM	Turkmenistan
BF	Burkina Faso	GR	Greece		Republic of Macedonia	TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	zw	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's	NZ	New Zealand	2	D040 # 0
CM	Cameroon		Republic of Korea	PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

Inte. onal Application No PCT/EP 98/06977

A. CLASSIFICATION OF SUBJECT MATTER IPC 6 C12N15/82 C12N C12N15/29 CO7K14/415 C12N15/10 C12N5/04 C12N5/14 A01H5/00 According to International Patent Classification (IPC) or to both national classification and IPC **B. FIELDS SEARCHED** Minimum documentation searched (classification system followed by classification symbols) IPC 6 C12N A01H Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched Electronic data base consulted during the international search (name of data base and, where practical, search terms used) C. DOCUMENTS CONSIDERED TO BE RELEVANT Citation of document, with indication, where appropriate, of the relevant passages Relevant to claim No. WO 95 15381 A (CHAPELLE ALBERT DE ;UNIV 1-5. JOHNS HOPKINS (US)) 8 June 1995 33 - 35see the whole document Α WO 97 37011 A (SETRATECH S A R L ; BORTS 1 - 35RHONA HARRIET (GB); LOUIS EDWARD JOHN (GB) 9 October 1997 see abstract see the whole document Α WO 90 07576 A (SETRATECH) 12 July 1990 1,7-27,31,32 see the whole document and specially page 5, line 16-31, examples 3-4 Further documents are listed in the continuation of box C. Patent family members are listed in annex. Special categories of cited documents: "T" later document published after the international filing date or priority date and not in conflict with the application but "A" document defining the general state of the lart which is not cited to understand the principle or theory underlying the considered to be of particular relevance invention "E" earlier document but published on or after the international "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another "Y" document of particular relevance; the claimed invention citation or other special reason (as specified) cannot be considered to involve an inventive step when the document is combined with one or more other such document, such combination being obvious to a person skilled in the art. "O" document referring to an oral disclosure, use, exhibition or other means document published prior to the international filing date but later than the priority date claimed "&" document member of the same patent family Date of the actual completion of the international search Date of mailing of the international search report 16 April 1999 03/05/1999 Name and mailing address of the ISA Authorized officer European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Tx. 31 651 epo ni, Mateo Rosell, A.M. Fax: (+31-70) 340-3016

Inter onal Application No
PCT/EP 98/06977

0.1011-		PCT/EP 98/06977
C.(Continu	ation) DOCUMENTS CONSIDERED TO BE RELEVANT	
	Citation of document, with indication where appropriate, of the relevant passages	Relevant to claim No.
Α	WO 97 01634 A (ANGELETTI P IST RICHERCHE BIO ;JIRICNY JOSEF (IT); PALOMBO FABIO () 16 January 1997 see page 1, line 18 - page 2, line 28 see page 58	1,2
A	PROLLA T A ET AL: "MLH1, PMS1, AND MSH2 INTERACTIONS DURING THE INITATION OF DNA MISMATCH REPAIR IN YEAST" SCIENCE, vol. 265, 19 August 1994, pages 1091-1093, XP000676403 cited in the application	1,2,4,5
A	A. WATANABE ET AL.,: "Genomic organization and expresssion of the human MSH3 gene" GENOMICS, vol. 31, 1996, pages 311-318, XP002099967 see the whole document and specially Fig.3	1-6
A	ACHARYA S ET AL: "hMSH2 forms specific mispair-binding complexes with hMSH3 and hMSH6" PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES OF USA, no. 93, October 1996, pages 13629- 13629, XP002080159 cited in the application sequences from this paper deposited in GenBank under AC= U61981 (hMSH3), and U54777 and U73732-7 (hMSH6). see the whole document	1-6
	K. LIU ET AL., : "characterization of the mouse Rep-3 gene: sequence similarities to bacterial and yeast mismatch-repair proteins" GENE, vol. 147, 1994, pages 169-177, XP002099968 see the whole document and specially Figure 2.	1-6
	I. IACCARINO ET AL.,: "MSH6, a Saccharomyces cerevisiae protein that binds to mismatches as a heterodimer with MSH2" CURRENT BIOLOGY, vol. 6, no. 4, April 1996, pages 484-486, XP002099969 sequence is deposited at GenBank under AC= Z47746. see the whole document	1-6

Inte. | onal Application No PCT/EP 98/06977

CICartin		PCT/EP 98/06977
	ation) DOCUMENTS CONSIDERED TO BE RELEVANT	
Category '	Citation of document, with indication where appropriate, of the relevant passages	Relevant to claim No.
A	CORRADI A ET AL: "CDNA SEQUENCE, MAP, AND EXPRESSION OF THE MURINE HOMOLOG OF GTBP, ADNA MISMATCH REPAIR GENE" GENOMICS, vol. 36, no. 2, 1 September 1996, pages 288-295, XP000613761 see the whole document and specially Fig.1	1-6
Ρ,Χ	DATABASE EMBL NUCLEOTIDE AND PROTEIN SEQUENCES,13 July 1998, XP002099368 HINXTON, GB AC= AJ007791. Arabidopsis thaliana mRNA for mismatch repair protein (MSH3) see abstract	1-6
Ρ,Χ	DATABASE EMBL NUCLEOTIDE AND PROTEIN SEQUENCES,1 August 1998, XP002099369 HINXTON, GB AC= 065607. Arabidopsis thaliana. Putative mismacht DNA repair protein see abstract	1-5
P,X	DATABASE EMBL NUCLEOTIDE AND PROTEIN SEQUENCES,16 October 1997, XP002099371 HINXTON, GB AC= AF009657. Arabidopsis thalian MutS homolog 6-like protein mRNA. see abstract -& K.M. CULLIGAN AND J.B. HAYS: "DNA mismatch repair in plants" PLANT PHYSIOLOGY, vol. 115, 1997, pages 833-839, XP002099372 see the whole document	1-6
-	DATABASE EMBL NUCLEOTIDE AND PROTEIN SEQUENCES,17 December 1998, XP002099373 HINXTON, GB AC= AJ131669. Triticum aestivum mRNA for MSH3 protein, MSH3 gene. see abstract	1-6
	DATABASE EMBL NUCLEOTIDE AND PROTEIN SEQUENCES, 12 October 1998, XP002099370 HINXTON, GB AC= AJ007792. Arabidopsis thaliana DNA mismatch repair protein, MSH6 gene. see abstract	1-6

information on patent family members

Inter onal Application No
PCT/EP 98/06977

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9515381	A 08-06-1995	EP 0730648 A JP 9506509 T US 5693470 A US 5837443 A	11-09-1996 30-06-1997 02-12-1997 17-11-1998
WO 9737011	A 09-10-1997	NONE	
WO 9007576	A 12-07-1990	FR 2641793 A AT 127519 T AU 4834390 A CA 2006549 A DE 68924174 D DE 68924174 T EP 0449923 A ES 2077058 T IE 72469 B JP 4503601 T	20-07-1990 15-09-1995 01-08-1990 26-06-1990 12-10-1995 14-03-1996 09-10-1991 16-11-1995 09-04-1997 02-07-1992
WO 9701634	A 16-01-1997	IT RM950434 A AU 6241296 A	27-12-1996 30-01-1997